arXiv:0809.4741v1 [math.PR] 27 Sep 2008

LARGE DEVIATIONS FOR THE LEAVES IN SOME RANDOM TREES

WLODEK BRYC, DAVID MINDA, AND SUNDER SETHURAMAN

Abstract. Large deviation principles and related results are given for a class of Markov chains associated to the "leaves" in random recursive trees and preferential attachment random graphs, as well as the "cherries" in Yule trees. In particular, the method of proof, combining analytic and Dupuis-Ellis type path arguments, allows for an explicit computation of the large deviation pressure.

1. Introduction and results

We consider in this article large deviations and related laws of large numbers and central limit theorems for a class of Markov chains associated to the number of leaves, or nodes of degree one, in preferential attachment random graphs and random recursive trees, and also the number of "cherries," or pairs of leaves with a common parent, in Yule trees. The random graphs studied model various networks such as pyramid schemes, chemical polymerization, the internet, social structures, genealogical families, among others. In particular, the leaf and cherry counts in these models are of interest, and have concrete interpretations. In Subsection 1.2, we discuss applications with these models and literature.

Define the nondecreasing Markov chain $\{Z_n : n \geq 1\}$ starting from initial state $Z_1 = k_0 \geq 0$, by its one-step transitions, for $n \geq 1$,

$$\Pr\left(Z_{n+1} - Z_n = v | Z_n\right) \ = \ \begin{cases} 1 - \frac{Z_n}{s_n} & \text{if } v = 1 \\ \frac{Z_n}{s_n} & \text{if } v = 0 \end{cases} \tag{1.1}$$

where $\{s_n : n \geq 1\}$ is a sequence of positive numbers such that

$$s_n \geq k_0 + n - 1, \text{ and } \frac{s_n}{n} \to \alpha \text{ for } 1 < \alpha < \infty \tag{1.2}$$

with convention $0/0 = 0$. Additionally, we also consider two special sequences, $s_n = n$ and $s_n = n/2$ with $k_0 \leq s_1$, related to some applications.

We first note, in this form, Z_n can be seen to represent the "mass" of red balls in a Polya-Eggenberger-type urn of two colors, red and blue, not necessarily tenable, where at each time n, proportional to the red balls mass, a (signed) mass $s_{n+1} - s_n$ of blue balls is deposited, otherwise one red ball and mass $s_{n+1} - s_n - 1$ of blue balls is added.

Date: September 26, 2008.

2000 *Mathematics Subject Classification.* 60F10;05C80.

Key words and phrases. large deviation, central limit, preferential attachment, planar oriented, uniformly random trees, leaves, cherries, Yule, random Stirling permutations.

Research supported in part by Taft Research Seminar 2006/07 and 2008/09.

Research supported in part by Taft Research Seminar 2008/09.

Research supported in part by NSF grant DMS-0504193.

Also, when $s_n = (k_0 + 2(n-1) + n\beta)/(1+\beta)$ and $\alpha = (2+\beta)/(1+\beta)$ for $\beta > -1$, the chain has interpretation as the number of leaves in preferential attachment random graphs with weight function $f(k) = k + \beta$. Later, in Subsection 1.2, we also remark that s_n can be taken as a random sequence with respect to the number of "generalized" leaves, or "buds," that is those nodes, possibly with degree greater than one, which however connect to only one other vertex, in preferential attachment graphs with random edge additions.

In addition, when $s_n = \alpha n$ for $\alpha = 1, 2$, Z_n is also the number of leaves in uniformly and planar oriented recursive trees respectively. In the latter case $\alpha = 2$, Z_n is also the number of "plateaux" in a random Stirling permutation of length $2n$.

Moreover, when $s_n = n/2$, Z_n can be seen as the number of cherries in Yule trees.

As the urns add mass of the opposite color, one should expect almost sure limits for the mean behavior and Normal fluctuations. In fact, by mostly martingale, combinatorial and urn methods, laws of large numbers (LLN) and central limit theorems (CLT) have been proved, at least in the examples mentioned above when s_n is linear with slope α (cf. Theorem 1.4). See [9, 36, 40] with respect to preferential attachment, [28], [41] with respect to recursive trees, [34] with respect to Yule trees, and [33] with respect to urns when α is a positive integer.

Characterizing the associated large deviations is a natural problem which gives insight into the properties of rare events, and seems less studied in urns or random graphs. Previous work has concentrated on analytic methods with respect to "subtraction" urn models–not applicable in our general setting [24]–or extensions of the Dupuis-Ellis weak convergence approach (cf. [20]) to different allocation models than ours [21], [44]. We note also some exponential bounds via martingale concentration inequalities are found in the case s_n is linear with slope α [15]. See also [4, 6, 12, 17, 19, 27] for other types of large deviations work in various random tree models.

In this context, our main results are to prove a large deviation principle (LDP) for Z_n/n with an explicitly computed "pressure," or Legendre transform of the associated rate function (Theorem 1.1). This is done in two different ways for the important case s_n is linear with $\alpha = 2$. Such explicit computations are not commonplace, and our "ODE" method is quite different from the methods in [24] where a quasi-linear PDE is solved, or in [44] where a finite-dimensional minimization problem is obtained.

Perhaps a main feature of our work is that the method given appears robust and applicable in diverse, not necessarily urn settings. In particular, we show the LDP for Z_n/n does not feel the disorder in the sequence s_n, is not dependent on the initial value Z_1, and is the same as for the chain with a regular, linear s_n with slope α. We mention that this is a consequence of a large deviation principle for the path interpolation of $Z_{\lfloor nt \rfloor}/n$ (Theorem 1.3), perhaps of interest in itself, that we establish, by the Dupuis-Ellis weak convergence approach.

In addition, aside from laws of large numbers which are trivially obtained, we prove a central limit theorem for Z_n through complex variables arguments with the pressure (Theorem 1.4). These alternate proofs of the LLN and CLT, although indirect, apply when the mass additions take on some negative non-integer values where less is known in the literature. Moreover, the results give a "quenched"

LLN, CLT and LDP's with respect to "generalized" leaves or buds in a preferential attachment scheme with random edge additions (see Subsection 1.2).

Our technique to prove the LDP for Z_n/n is to consider the recurrence relation for $m_n(\lambda) = E[\exp\{\lambda Z_n\}]$ obtained from (1.1):

$$m_{n+1}(\lambda) \;=\; (1-e^{\lambda})\frac{m_n'(\lambda)}{s_n} + e^{\lambda}m_n(\lambda). \tag{1.3}$$

Dividing through by $m_n(\lambda)$, we write

$$\frac{m_{n+1}(\lambda)}{m_n(\lambda)} \;=\; \frac{1-e^{\lambda}}{s_n/n}\frac{m_n'(\lambda)}{nm_n(\lambda)} + e^{\lambda}. \tag{1.4}$$

The idea now is to take the limit on n in the above display. When the "pressure" Λ exists, it satisfies $\Lambda(\lambda) = \lim_{n\to\infty}(1/n)\log m_n(\lambda)$. In this case, it is natural to suppose that the limits

$$\Lambda'(\lambda) \;=\; \lim_{n\to\infty}\frac{m_n'(\lambda)}{nm_n(\lambda)} \tag{1.5}$$

$$e^{\Lambda(\lambda)} \;=\; \lim_{n\to\infty}\frac{m_{n+1}(\lambda)}{m_n(\lambda)} \tag{1.6}$$

both exist. Then, from (1.4), we can write the ODE

$$e^{\Lambda(\lambda)} \;=\; \frac{1-e^{\lambda}}{\alpha}\Lambda'(\lambda) + e^{\lambda}; \quad \Lambda(0) = 0. \tag{1.7}$$

One can compute the solution of this differential equation (cf. (1.12)) and show it is unique.

The main task is to show that the pressure and limits (1.5) and (1.6) exist. But, the pressure exists as a consequence of the path LDP for $Z_{\lfloor nt\rfloor}/n$ by contraction principle. We note, in principle, one can try to compute the pressure or the rate function from (1.13) by the calculus of variations, but we found it difficult to solve the associated Euler equations (cf. near (4.1)).

Finally, we show (1.5) and (1.6) exist by extending $m_n(\lambda)$ to the complex plane, and then analyzing its zeroes and analytic properties. These estimates are also useful for the central limit theorem arguments.

We now mention a different approach, in the spirit of [24], when s_n is linear with slope $\alpha = 2$ and also interestingly $\alpha = 1/2, 1$, to compute the pressure from analysis of the generating function $G(\lambda, z) = \sum_{n\geq 1} m_n(\lambda)z^{n-1}$. From (1.3), we can write the linear PDE

$$\frac{\partial G}{\partial z}(1-e^{\lambda}z) + \frac{e^{\lambda}-1}{\alpha}\frac{\partial G}{\partial \lambda} \;=\; e^{\lambda}G. \tag{1.8}$$

One can solve implicitly this PDE, and locate at least heuristically a singular point. Then, formally, from root test asymptotics, the pressure would be the reciprocal of the location of the singularity.

The difficulty is in establishing the analyticity of the solution and identifying its singularity. For urn models of "subtraction type" where the mass added is an integer, and which satisfies a balance condition, [24] uses this program to obtain large deviations and the CLT. However, the cases s_n is linear with slope $\alpha = 1/2, 1, 2$, and more generally the urns associated with non-integer s_n are not covered by their arguments which seem to rely on integer additions with a certain "negative" structure. On the other hand, we are able to supply the needed analyticity and singularity identification when s_n has slopes $\alpha = 1/2, 1, 2$, and in this way prove the LDP for Z_n/n (Theorem 1.1) in these cases.

The plan of the paper is to state the results in Subsection 1.1, discuss applications to random graphs in Subsection 1.2, give the generating function proof of Theorem 1.1 in Section 2, prove the path LDP (Theorem 1.3) in Section 3, give the ODE-method proof of Theorem 1.1 and prove the LLN and CLT (Theorem 1.4) in Section 4, and conclude in Section 5.

1.1. **Results.** We recall the setting for large deviations. A sequence $\{X_n\}$ of random variables with values in a separable complete metric space $\mathbf{X}$ satisfies the large deviation principle (LDP) with speed n and rate function $I : \mathbf{X} \to [0, \infty]$ if I has compact level sets $\{\mathbf{x} : I(\mathbf{x}) \le a\}$, and for every Borel set $U \in \mathcal{B}_{\mathbf{X}}$,

$$\begin{aligned} -\inf_{\mathbf{x}\in U^\circ} I(\mathbf{x}) &\le \liminf_{n\to\infty} \frac{1}{n} \log \Pr(X_n \in U) \\ &\le \limsup_{n\to\infty} \frac{1}{n} \log \Pr(X_n \in U) \le -\inf_{\mathbf{x}\in \bar{U}} I(\mathbf{x}). \end{aligned} \tag{1.9}$$

[Here U° is the interior of U and $\bar{U}$ is the closure of U.]

Often the rate function is given in terms of the Legendre transform of the "pressure" when it exists. When $\mathbf{X} = \mathbb{R}$, this representation takes the form

$$I(x) = \sup_{\lambda\in\mathbb{R}} \big\{\lambda x - \log \Lambda(\lambda)\big\}, \tag{1.10}$$

where we recall the pressure satisfies

$$\Lambda(\lambda) := \lim_{n\to\infty} \frac{1}{n} \log E[e^{\lambda Z_n}]. \tag{1.11}$$

Recall now the Markov chain Z_n (1.1) corresponding to sequence $\{s_n\}$ and parameter $1 < \alpha < \infty$ (1.2).

Theorem 1.1. *The sequence Z_n/n satisfies LDP with speed n and good strictly convex rate function I given by* (1.10) *with pressure*

$$\Lambda(\lambda) = -\log\left(\frac{\alpha}{e^\lambda - 1} \int_0^\lambda \left(\frac{e^s - 1}{e^\lambda - 1}\right)^{\alpha-1} ds\right) \qquad \text{for } \lambda \ne 0 \tag{1.12}$$

and $\Lambda(0) = 0$.

Remark 1.2. We note, when $s_n = \alpha n$ with $\alpha = 1/2, 1$ and $k_0 \le s_1$, the LDP is also found, with pressure given by formula (1.12), using the generating function approach (cf. Section 2). For $\alpha = 1/2$ and integer $\alpha \ge 1$, the integral in (1.12) can be evaluated explicitly.

We now consider the LDP for the family of stochastic processes $\{X_n(t) : 0 \le t \le 1\}$ obtained by linear interpolation of the Markov chain (1.1),

$$X_n(t) := \frac{1}{n} Z_{\lfloor nt\rfloor - k_0 + 1} + \frac{nt - \lfloor nt\rfloor}{n} (Z_{\lfloor nt\rfloor - k_0 + 2} - Z_{\lfloor nt\rfloor - k_0 + 1}) \ \text{ for } t \ge \frac{k_0}{n}$$

and $X_n(t) := t$ for $0 \le t \le k_0/n$. The trajectories of $X_n(t)$ are non-decreasing Lipschitz functions with constant at most 1.

Theorem 1.3. *As a sequence of* $C([0,1];\mathbb{R})$*-valued random variables,* X_n *satisfies the LDP with the rate function* $I : C([0,1];\mathbb{R}) \to [0,\infty]$ *given by*

$$I(\varphi) \;=\; \int_0^1 \left[\dot{\varphi}(t)\log\frac{\alpha t\dot{\varphi}(t)}{\alpha t-\varphi(t)} + (1-\dot{\varphi}(t))\log\frac{\alpha t(1-\dot{\varphi}(t))}{\varphi(t)}\right] dt \tag{1.13}$$

if $\varphi(t)$ *is differentiable for almost all* t*,* $\varphi(0) = 0$*,* $0 \le \dot{\varphi} \le 1$*, and the integral converges; otherwise,* $I(\varphi) = \infty$*.*

By the contraction principle, Theorem 1.3 implies the LDP for Z_n/n with the rate function given by the variational expression

$$I(x) \;=\; \inf\Big\{I(\varphi) :\; \varphi(0) = 0,\; \varphi(1) = x\Big\}. \tag{1.14}$$

In general, optimal trajectories are not straight lines—exceptions are the LLN trajectory $\varphi_\alpha(t) = t\alpha/(\alpha+1)$ and the extreme case $\varphi(t) = t$—but they try to stay near the LLN line (for which $I(\varphi_\alpha) = 0$) to minimize cost before going to destination x (cf. Fig. 1).



Figure 1. Thick curves are numerical solutions of the Euler equations for (1.14) with $\alpha = 2$ for $x = 0.13, 2/3, 0.85, 1$. Dashed lines are straight lines from $(0,0)$ to $(1,x)$.

Lemmas for the proof of Theorem 1.1 give Normal approximation. The law of large numbers also follows from Theorem 1.1.

Theorem 1.4. *We have*

$$\frac{Z_n}{n} \;\xrightarrow{\text{a.s.}}\; \frac{\alpha}{\alpha+1}$$

and also

$$\frac{1}{\sqrt{n}}(Z_n - E[Z_n]) \;\xrightarrow{\text{D}}\; N(0,\sigma^2) \quad\text{where}\quad \sigma^2 \;=\; \frac{\alpha^2}{(1+\alpha)^2(2+\alpha)}.$$

Remark 1.5. The conclusions of Theorem 1.4 also hold when $s_n = \alpha n$ with $\alpha = 1/2, 1$, and $k_0 \le s_1$ (cf. Subsection 4.3).

1.2. **Applications to random graph models.** With respect to random graphs, the Markov chain Z_n, representing the number of leaves, sits in the intersection of at least two models, that is preferential attachment graphs with linear-type weights, and uniformly and planar oriented trees. Also, when $s_n = \alpha n$ for $\alpha = 1/2$, Z_n represents the count of cherries in Yule trees.

1.2.1. *Preferential attachment graphs.* Preferential attachment graphs have a long history dating back to Yule (cf. [35]). However, since the work of Barabasi-Albert [1, 5], these graphs have been of recent interest with respect to modeling of various "real-world" networks such as the internet (WWW), and social and biological communities. Leaves, or nodes with degree one, in these networks of course represent sites with one link, or members at the "fringe." [See books [23], [15], [11] for more discussion.]

The idea is to start with an initial connected graph G_1 with a finite number of vertices, and say no self-loops (so that the vertices have well-defined degrees). At step 1, add another vertex, and connect it to a vertex x of G_1 preferentially, that is with probability proportional to its weight, $f(d_x)/\sum_{y\in G_1} f(d_y)$, to form a new graph G_2. Continue in this way by adding a new vertex and connecting it preferentially to form G_k for $k \geq 1$. Here, the "weight" of a vertex is a function f of its degree d_x. When $f : \mathbb{N} \to \mathcal{R}_+$ is increasing, already well connected vertices tend to become better connected, a sort of reinforcing effect.

Our results are applicable to linear weights, $f(k) = k + \beta$, for $\beta > -1$, which correspond to certain "power law" mean degree sequences. Namely, let $Z_k(n)$ be the number of vertices in G_n with degree k for $k \geq 1$. It was shown, by martingale arguments, in [9], [36], and by embedding into branching processes, in [40] that $Z_k(n)/n \to r_k$ a.s. where

$$r_k \ = \ \frac{2+\beta}{k+\beta}\prod_{j=1}^{k}\frac{j+\beta}{j+2+2\beta} \ \sim \ 1/k^{3+\beta}.$$

From the applications point of view, the parameter β can sometimes be matched to empirical network data where similar power-law behavior is observed.

As the number of vertices with degree 1, or the "leaves" $Z_1(n)$, increases by one in the next step when a non-leaf is selected, and remains the same when a leaf is chosen, we see that $Z_1(n)$ corresponds to the Markov chain Z_n with s_n as specified below. Since at each step the total degree of the graph augments by two, the probability at step n that a vertex $x \in G_n$ is selected is $(d_x+\beta)/(d_{G_1}+2(n-1)+(n-1+|G_1|)\beta)$. Then, $s_n = \big(d_{G_1} + 2(n-1) + (n-1+|G_1|)\beta\big)/(1+\beta)$ and $\alpha = (2+\beta)/(1+\beta)$. Here d_{G_1} and $|G_1|$ are the total degree and number of vertices in G_1 respectively.

One can also "randomize" the model by adding a random number of edges at each step: Let $\{\gamma_i\}$ be a sequence of independent identically distributed random variables on $\mathbb{N}$ with finite mean, $\bar\gamma = E[\gamma_1] < \infty$. Then, at step n, we add a new vertex to the graph G_n and connect it to a node selected preferentially from G_n with γ_n edges put between them. The effect of these random edge additions is to randomize further the weight given the nodes in the graph; the "deterministic" model above is when $P(\gamma_1 = 1) \equiv 1$. In [3], by embedding into branching processes, it was shown, when $E(\gamma_1 \log \gamma_1) < \infty$, that $Z_k(n)/n \xrightarrow{P} r_k$ where r_k is given by an integral formula with asymptotics $r_k \sim 1/k^{3+\beta/\bar\gamma}$. See [16] for an even more general randomized model where also a LLN is proved. Other models with different random edge addition schemes are found in [19], [31].

In this "randomized" model, we now define the notion of "generalized leaves" or "buds," that is, those nodes which connect to exactly one other vertex, albeit possibly with several edges linking them. Leaves are those buds with exactly one edge connection. Similar to the leaves in the deterministic setting, at step n, the bud count increases by one exactly when the new vertex, a fortiori a bud, connects to a non-bud in G_n, and remains the same when the new vertex links to a bud in G_n.

Then, the Markov chain Z_n, representing the number of buds at step n, corresponds to $s_1 = (d_{G_1} + |G_1|\beta)/(1+\beta)$ and for $n \geq 2$,

$$s_n \;=\; \frac{1}{1+\beta}\Big[d_{G_1} + 2\sum_{i=1}^{n-1}\gamma_i + \big(n-1+|G_1|\big)\beta\Big],$$

which satisfies assumption (1.2), by the strong law of large numbers, with $\alpha = (2\bar{\gamma}+\beta)/(1+\beta)$ a.s. In particular, by our results, we obtain "quenched" LLN, CLT and LDP's, with probability one with respect to $\{\gamma_i\}$, for the buds in this scheme.

1.2.2. *Random recursive trees.* Random recursive trees are also well-established models, dating to the 1960's, with applications to data sorting and searching, pyramid schemes, spread of epidemics, chemical polymerization, family trees (stemma) of copies of ancient manuscripts etc. Leaves in these trees correspond to "shutouts" with respect to pyramid schemes, nodes with small "affinity" in polymerization models, "terminal copies" in stemma of manuscripts etc. See [41], [42] and references therein (e.g. [37] etc.) for more discussion. Below, we mention connections with Stirling permutations.

Similar to preferential attachment, the recursive schemes form a sequence of trees. One starts with a single vertex labeled 0, and then adds a vertex at step $n \geq 1$, labeled n, to one of the n nodes already present. When the choice is made uniformly, the model forms a *uniformly* recursive tree. However, when the choice is made proportional to the degree, a non-uniform or *plane oriented* tree is grown. The interpretation is that in uniform trees, at each distance from the root 0, there is no ordering of the labels. However, in plane-oriented trees, different orders give rise to distinct trees; that is, after selecting a parent node at random at step n with k children with a certain cyclic order of labels, there are $k+1$ locations where the new label n can be inserted.

Plane oriented trees are similar to preferential attachment graphs with $\beta = 0$ (cf. Chapter 4 [23]), and the number of leaves corresponds to the Markov chain with $k_0 = 1$, $s_n = 2n-1$ for $n \geq 1$, and $\alpha = 2$. On the other hand, the number of leaves in uniformly recursive trees corresponds to the case $k_0 = 1$, $s_n = n$ for $n \geq 1$, and $\alpha = 1$. With respect to both types of recursive trees, LLN's and CLT's have also been proved by combinatorial, urn and martingale methods (see [41], [28]). So, in this context, our results give alternate proofs of the LLN and CLT for these recursive trees, and also an LDP for Z_n/n. We note also Theorem 1.3 yields a path LDP for planar oriented trees.

We comment now on recent connections of planar oriented trees with Stirling permutations (cf. [29], [30]). A Stirling permutation of length $2n$ is a permutation of the multiset $\{1,1,2,2,\ldots,n,n\}$ such that for each $i \leq n$ the elements occurring between the two i's are larger than i (cf. [26]). It turns out that each permutation is a distinct code for a plane oriented recursive tree with $n+1$ vertices.

Quoting from [29], consider the depth first walk which starts at the root, and goes first to the leftmost daughter of the root, explores that branch (recursively, using the same rules), returns to the root, and continues to the next daughter and so on. Each edge is passed twice in the walk, once in each direction. Label the edges in the tree according to the order in which they were added–edge j is added at step j and connects vertex j to an previously labeled vertex. The plane recursive tree is coded by the sequence of labels passed by the depth first walk. With respect to a tree with $n+1$ vertices, the code is of length $2n$, where each of the labels $1, 2, \dots, n$ appears twice. Adding a new vertex means inserting the pair $(n+1)(n+1)$ in the code in one of the $2n+1$ places.

In a Stirling permutation $a_1 a_2 \cdots a_{2n}$, the index $0 \le i \le 2n$ is a plateau if $a_i = a_{i+1}$ (where $a_{2n+1} = 0$). [29] shows that the number of leaves in a plane oriented tree with $n+1$ vertices is the number of plateaux in a random Stirling permutation. See [29] for more details.

1.2.3. *Yule trees.* Since Yule's influential 1924 paper [43], Yule trees, among other processes, have been used widely to model phylogenetic evolutionary relationships between species (see [2] for an interesting essay). In particular, the counts of various shapes and features of these trees can be studied, and matched to empirical data to test evolutionary hypotheses. In [34], a LLN and CLT is proved for the number of cherries, or pairs of leaves with a common parent, in Yule trees. Associated confidence intervals are computed, and some empirical data sets are examined to see their compatibility with "Yule tree" genealogies. Other related statistical tests and limit results can be found in [7, 8, 39].

In the Yule tree process, one starts with a root vertex. It will split into two daughter nodes at step 2, each of which is equally likely to split into two children at step 3. At step n, one of the n leaves in the tree is chosen at random, and it then splits into two daughters, and so on. It is easy to see that the number of cherries at step n is given by the Markov chain Z_n with $s_n = n/2$ and $k_0 = 0$. Correspondingly, our results give different proofs of the LLN and CLT for the cherry counts, as well as an LDP for Z_n/n.

2. PROOF OF THEOREM 1.1: CASES $s_n = \alpha n$, $\alpha = 1/2, 1, 2$

In this section we give a proof of the LDP in Theorem 1.1, via an analysis of a generating function, for the special case $s_n = \alpha n$ for $\alpha = 2$, at the intersection of several models, and two additional values $\alpha = 1/2, 1$ not covered by Theorem 1.3.

The LDP is obtained by Gärtner-Ellis theorem [18, Theorem 2.3.6] from the following.

Proposition 2.1. *Suppose $s_n = \alpha n$ with $\alpha = 1/2$, 1, or 2, and $Z_1 = k_0 \le s_1$. The limit* (1.11) *exists, and is given by the smooth function*

$$\begin{aligned}
\Lambda(\lambda) \quad &= \quad \begin{cases} \log\left(\frac{\sqrt{e^\lambda - 1}}{\arctan\left(\sqrt{e^\lambda-1}\right)}\right) & \text{if } \lambda > 0 \\ \log\left(\frac{\sqrt{1-e^\lambda}}{\operatorname{arctanh}\left(\sqrt{1-e^\lambda}\right)}\right) & \text{if } \lambda < 0 \end{cases} && \text{for } \alpha = 1/2 \\
&= \quad \log \frac{e^\lambda - 1}{\lambda} && \text{for } \alpha = 1 \\
&= \quad \log\left(\frac{\left(e^\lambda - 1\right)^2}{2\left(e^\lambda - 1 - \lambda\right)}\right) && \text{for } \alpha = 2 \qquad (2.1)
\end{aligned}$$

when $\lambda \neq 0$, and $\Lambda(0) = 0$.

We follow analytic arguments adapted from [24]. Since $0 \le k_0 \le Z_n \le n+k_0-1$, we get $0 < m_n(\lambda) \le e^{(n+k_0)\lambda^+}$ with $\lambda^+ = \max\{\lambda, 0\}$. Therefore, for all complex z

with $|z| < e^{-\lambda^+}$, $G(z, \lambda)$ is well defined and satisfies (1.8) with the initial condition $G(0, \lambda) = e^{k_0 \lambda}$. The coefficients of this PDE do not vanish in the regions

$$\mathcal{D}_+ = \big\{(z, \lambda) : \lambda > 0, |z| < e^{-\lambda}\big\},\ \mathcal{D}_- = \big\{(z, \lambda) : \lambda < 0, |z| < 1\big\}.$$

For $\lambda \neq 0$, the PDE can be solved by the method of characteristics. Clearly, $m_n(\lambda)$ are the coefficients in the Taylor expansion at $z = 0$ of the solution, and $e^{-\Lambda(\lambda)}$ is the radius of convergence of the series that can be determined by singularity analysis.

For $\alpha = 1$, $k_0 = 0, 1$; using initial condition $G(0, \lambda) = e^{k_0 \lambda}$ we get

$$G(\lambda, z) \;=\; e^{k_0(z(e^\lambda - 1) - \lambda)} \frac{e^\lambda - 1}{1 - \exp(z(e^\lambda - 1) - \lambda)}.$$

Hence, the singularity of G as a function of z nearest to the origin is a simple pole at

$$z_0 \;=\; \frac{\lambda}{e^\lambda - 1}.$$

By Darboux's asymptotic method [38, Ch. 8],

$$\frac{1}{n} \log E(\exp(\lambda Z_n)) \;\to\; \log \frac{e^\lambda - 1}{\lambda}$$

and the LDP follows.

Next, consider $\alpha = 1/2$. In this case, the solutions of the PDE differ depending on the region $\mathcal{D}_\pm$, but are explicit so there are no difficulties in constructing their analytic extensions. Using the initial condition $k_0 = 0$, we have $G(0, \lambda) = 1$, and the solution of (1.8) is

$$G(\lambda, z) \;=\; \begin{cases} \dfrac{\sqrt{e^\lambda - 1}}{\tan\big(\arctan\sqrt{e^\lambda - 1} - z\sqrt{e^\lambda - 1}\big)} & \lambda > 0 \\ \dfrac{\sqrt{1 - e^\lambda}}{\tanh\big(\operatorname{arctanh}\sqrt{1 - e^\lambda} - z\sqrt{1 - e^\lambda}\big)} & \lambda < 0 \end{cases}$$

Hence, the singularity of G as a function of z nearest to the origin is a simple pole at

$$z_0 \;=\; \begin{cases} \dfrac{\arctan\sqrt{e^\lambda - 1}}{\sqrt{e^\lambda - 1}} & \lambda > 0 \\ \dfrac{\operatorname{arctanh}\sqrt{1 - e^\lambda}}{\sqrt{1 - e^\lambda}} & \lambda < 0 \end{cases}$$

Once again, by Darboux's asymptotic method [38, Ch. 8],

$$\frac{1}{n} \log E(\exp(\lambda Z_n)) \;\to\; \log 1/z_0$$

and the LDP follows.

For $\alpha = 2$, the method of characteristics gives the following answer.

Lemma 2.2. *Suppose φ is a function of one complex variable, analytic in a domain $\mathcal{D}$ containing $(-\infty, -2)$. Then*

$$G(z, \lambda) \;:=\; (e^\lambda - 1)^2 \varphi(z(e^\lambda - 1)^2 + 2\lambda - 2e^\lambda). \tag{2.2}$$

satisfies the PDE (1.8) *for all* $\lambda \neq 0$ *and* $|z|$ *small enough. Furthermore, the initial condition is fulfilled at* $\lambda \neq 0$ *if*

$$\varphi(2\lambda - 2e^{\lambda}) \;=\; \frac{e^{k_0\lambda}}{(1-e^{\lambda})^2}. \tag{2.3}$$

Proof. The verification of the initial condition is trivial, and the verification of the PDE is a straightforward calculation. Denoting for conciseness $\psi(z,\lambda) = \varphi(z(e^{\lambda}-1)^2 + 2\lambda - 2e^{\lambda})$, $\psi^{(1)}(z,\lambda) = \varphi'(z(e^{\lambda}-1)^2 + 2\lambda - 2e^{\lambda})$ we verify, for $\lambda \neq 0$ and z such that $z(e^{\lambda}-1)^2 + 2\lambda - 2e^{\lambda} \in \mathcal{D}$, that

$$\frac{\partial G(z,\lambda)}{\partial \lambda} \;=\; 2(e^{\lambda}-1)e^{\lambda}\psi(z,\lambda) + 2(e^{\lambda}-1)^3(ze^{\lambda}-1)\psi^{(1)}(z,\lambda)$$

and

$$\frac{\partial G(z,\lambda)}{\partial z} \;=\; (e^{\lambda}-1)^4\psi^{(1)}(z,\lambda).$$

Equation (1.8) now follows by a calculation. □

Our next goal is to show that one can find a solution of (1.8) which can be analytically extended in variable z to a large enough domain. To this end, we need to analyze (2.3) with complex argument. The basic plan consists of noting that function $f(x) = 2(x - e^{x})$ is analytic, strictly decreasing for $x > 0$, strictly increasing for $x < 0$, and $f(0) = -2$. The derivative $f'(x)$ vanishes only at $x = 0$, so $f\big|_{[0,\infty)}$ and $f\big|_{(-\infty,0]}$ have continuous inverses $h_+ : (-\infty,-2] \to [0,\infty)$ and $h_- : (-\infty,-2] \to (-\infty,0]$, and both are analytic on $(-\infty,-2)$.

Clearly, if we define

$$\varphi_{\pm}(\lambda) \;=\; \frac{e^{k_0 h_{\pm}(\lambda)}}{(e^{h_{\pm}(\lambda)}-1)^2}\,, \tag{2.4}$$

then φ_+ satisfies (2.3) for $\lambda > 0$, and φ_- satisfies (2.3) for $\lambda < 0$. The goal is therefore to find the appropriate analytic extensions of the functions $h_{\pm}$.

2.1. Construction of an analytic extension.

We need to analyze $f(z) = 2(z - e^{z})$. The closely related function $z + e^{z}$ appears in [32, page 116] but proofs are not included there; we give details for $f(z) = 2(z - e^{z})$ for completeness.

The proof relies on the following univalence criterion.

Lemma 2.3 (Wolff-Warschawski-Nishiro)**.** *If* g *is holomorphic in a convex region* Ω *and* $g'(\Omega) \subset H$*, a half-plane with* $0 \in \partial H$*, then* g *is one-to-one on* Ω*.*

Proof. This is [22, Theorem 2.16 on page 47] applied to the function $e^{i\theta}g(z)$ with a real constant θ chosen appropriately to rotate H. □

Lemma 2.4. f *is a one-to-one mapping of the half-closed strip* $\Sigma = \{z : 0 < \Im(z) \leq \pi\}$ *onto a slit closed half-plane:* $\{w : \Im(w) \leq \pi\} \setminus (-\infty,-2]$. *The boundary correspondence is as follows:* f *maps* $\mathbb{R} + \pi i$ *injectively onto* $\mathbb{R} + 4\pi i$*, and* f *is one-to-one on both* $[0,+\infty)$ *and* $(-\infty,0]$ *and maps each onto* $(-\infty,-2]$ *(cf. Fig. 2).*

Proof. We write $f(z) = g(\zeta)$, where $\zeta = e^{z}$, $g(\zeta) = 2(\log\zeta - \zeta)$ and log denotes the principal branch of the logarithm. The function $\zeta = e^{z}$ is a one-to-one mapping on the strip Σ. The image of the interior of Σ is the upper half-plane $\mathbb{H} = \{\zeta : \Im(\zeta) > 0\}$. Furthermore, the upper edge $\mathbb{R} + \pi i$ of Σ is mapped onto $(-\infty,0)$ and the bottom edge $\mathbb{R}$ is carried onto $(0,+\infty)$.

The derivative of $g(\zeta)$ is $g'(\zeta) = 2\bar{\zeta}/|\zeta|^2 - 2$. In particular, $\Im(g'(\zeta)) = -2\Im(\zeta)/|\zeta|^2 < 0$ for $\zeta \in \mathbb{H}$. By Lemma 2.3 with $\Omega = \mathbb{H}$, g is one-to-one on the half-plane $\mathbb{H}$. Under g the image of $\mathbb{H}$ is $\{w : \Im(w) \leq 2\pi\} \setminus (-\infty, -2]$, and on the boundary we have $g(\mathbb{R}^-) = \mathbb{R} + 2\pi i$ and $g(\mathbb{R}^+)$ is the slit $(-\infty, -2]$ twice covered with $g(1) = -2$. Perhaps this latter statement is easier seen directly from f; the slit is covered twice: $f((-\infty, 0]) = f([0, +\infty)) = (-\infty, -2]$. $\square$



FIGURE 2. Composition of maps $f(z) = g(e^z)$ for the proof of Lemma 2.4.

We investigate the conformal mapping $f|\Sigma$ in more detail. The preimage of $[-2, \infty)$ under $f|\Sigma$ is the curve γ given by $\Re z = \log(\Im z/\sin \Im z)$, $0 \leq \Im z < \pi$. This curve begins at the origin and becomes asymptotic to $\mathbb{R} + \pi i$ in the positive direction. By removing the curve γ, Σ is cut into two parts, Σ_- and Σ_+. Σ_- is bounded by $\mathbb{R}^-$, γ and $\mathbb{R} + \pi i$ with the latter line part of Σ_-. The region Σ_+ is bounded by γ and $\mathbb{R}^+$. Then $f|\Sigma_-$ is a conformal mapping of Σ_- onto the half-closed strip $S = \{w : 0 < \Im w \leq 2\pi\}$ with $f(\mathbb{R} + \pi i) = \mathbb{R} + 2\pi i$, $f(\mathbb{R}^-) = (-\infty, -2)$ and $f(\gamma) = [-2, +\infty)$. Similarly, $f|\Sigma_+$ is a conformal mapping of Σ_+ onto the lower half-plane $\overline{\mathbb{H}} = \{\bar{z} : z \in \mathbb{H}\}$, where $\bar{z}$ denotes the complex conjugate of z, with $f(\mathbb{R}^+) = (-\infty, -2)$ and $f(\gamma) = [-2, +\infty)$. (See Fig 3.)

Both maps $f|\Sigma_-$ and $f|\Sigma_+$ extend to conformal maps of larger regions. Because $f(\bar{z}) = \overline{f(z)}$, f maps $\overline{\Sigma_+}$ conformally onto $\mathbb{H}$. As $f(\mathbb{R}^+) = (-\infty, -2)$, f is a conformal map of $\Omega_+ := \Sigma_+ \cup \mathbb{R}^+ \cup \overline{\Sigma_+}$ onto the slit plane $\bar{\mathbb{H}} \cup (-\infty, -2) \cup \mathbb{H} = \mathbb{C} \setminus [-2, +\infty)$. Let $h_+ : \mathbb{C} \setminus [-2, +\infty) \to \Omega_+$ be the inverse function for $f|\Omega_+$. The conformal extension of $f|\Sigma^-$ is more involved to describe. The fact that $f(\bar{z}) = \overline{f(z)}$ implies f maps $\overline{\Sigma_-}$ conformally onto $\bar{S}$. Since $f(\mathbb{R}^-) = (-\infty, -2)$, f is a conformal map of $\Delta_0 = \Sigma_- \cup \mathbb{R}^- \cup \overline{\Sigma_-}$ onto the slit closed strip $S_0 = S \cup (-\infty, -2) \cup \bar{S} = \{w : |\Im w| \leq 2\pi\} \setminus [-2, +\infty)$ with the upper (lower) edge of Δ_0 corresponding to the upper (lower) edge of S_0. It is straightforward to verify that $f(z + 2\pi i) =$



Figure 3. Conformal maps $f|\Sigma_-$ and $f|\Sigma_+$.

$f(z)+4\pi i$ for each $n \in \mathbb{Z}$. This functional relationship implies that f is a conformal map of $\Delta_n = \Delta_0 + 2\pi ni$ onto $S_n = S_0 + 4\pi ni$ for each $n \in \mathbb{Z}$. Hence, f is a conformal map of $\Omega_- := \cup_{n\in\mathbb{Z}}\Delta_n$ onto the infinitely slit plane $S_\infty := \bigcup_{n\in\mathbb{Z}} S_n = \mathbb{C} \setminus \bigcup_{n\in\mathbb{Z}} \big([-2,+\infty) + 4\pi ni\big)$. Let $h_- : S_\infty \to \Omega_-$ be the inverse function of $f|\Omega_-$. Because $\mathbb{C} \setminus [-2,+\infty) \supset S_\infty$, we may regard h_+ as defined on S_∞, so $h_\pm$ have a common domain. (See Fig. 4.)

The two conformal maps $h_\pm$ just constructed provide analytic extensions to S_∞ of the real-valued functions $h_\pm$. This allows us to define a pair of functions

$$\varphi_\pm(z) := \frac{e^{k_0 h_\pm(z)}}{(e^{h_\pm(z)} - 1)^2} \tag{2.5}$$

which are analytic in S_∞.

Lemma 2.5. *For each point $u \in (-\infty,-2)$ the power series expansion of $\varphi_\pm$ has radius of convergence $2-u$, $\varphi_\pm$ is analytic in the slit disk $D(u,r)\setminus[-2,+\infty)$, where $r = r(u) = \sqrt{(u+2)^2+16\pi^2}$. Furthermore, $\varphi_\pm(w) \approx -1/(w+2)$ as $w \to -2$ in S_∞.*

Proof. For each point $u \in (-\infty,-2)$ the power series expansion of $h_\pm$ has radius of convergence $|u+2|$, the distance from u to -2 because -2 is the closest singularity of $h_\pm$. Also, $h_\pm$ is analytic in the slit disk $D(u,r) \setminus [-2,+\infty)$, where $r = r(u) = \sqrt{(u+2)^2+16\pi^2}$ is the distance from u to $-2 \pm 4\pi i$.

Note that singularities of φ_- that arise from $h_-(w) = 2n\pi i$ are located at the slits taken out of S_∞ and that $h_+(w) \in \Omega_+$ cannot take values in $2n\pi i$. Thus $\varphi_\pm$ is also analytic in the slit disk $D(u,r) \setminus [-2,+\infty)$.



FIGURE 4. Conformal maps $f|\Omega_+$ and $f|\Omega_-$ with $\Omega_+ := \Sigma_+ \cup \mathbb{R}^+ \cup \overline{\Sigma_+}$ and $\Omega_- = \bigcup_{n=-\infty}^{\infty} \Delta_n$.

Substituting $w = 2(z - e^z)$, we see that

$$\begin{aligned}\lim_{w\to -2,\ w\in S_\infty} (w+2)\varphi_\pm(w) &= \lim_{w\to -2,\ w\in S_\infty} e^{k_0 h_\pm(w)} \frac{w+2}{(e^{h_\pm(w)}-1)^2} \\ &= 2 \lim_{z\to 0,\ z\in\Omega_\pm} e^{k_0 z} \frac{1+z-e^z}{(e^z-1)^2} = -1.\end{aligned}$$

□

Conclusion of proof of Proposition 2.1 for $\alpha = 2$. We now prove (1.11) and identify the limit. Using functions $\varphi_\pm$ constructed above, define

$$\widetilde{G}(z,\lambda) := \begin{cases} (e^\lambda-1)^2 \varphi_+(z(e^\lambda-1)^2 + 2\lambda - 2e^\lambda) & \text{if } \lambda > 0 \\ (e^\lambda-1)^2 \varphi_-(z(e^\lambda-1)^2 + 2\lambda - 2e^\lambda) & \text{if } \lambda < 0\ . \end{cases} \tag{2.6}$$

From Lemma 2.2 we see that function $\widetilde{G}(z,\lambda)$ satisfies (1.8) for all $(z,\lambda) \in \mathcal{D}_+ \cup \mathcal{D}_-$. By uniqueness of the PDE solution in each of the two regions, $G(z,\lambda) = \widetilde{G}(z,\lambda)$ for all $(z,\lambda) \in \mathcal{D}_+ \cup \mathcal{D}_-$. In particular, for each fixed $\lambda \neq 0$, $\limsup_{n\to\infty}(m_n(\lambda))^{1/n}$ is the reciprocal of the radius of convergence of the series expansion of $\widetilde{G}(z,\lambda)$ at $z = 0$. The latter is $(1-e^\lambda)^{-2}$ times the radius of convergence for $\varphi_{sign(\lambda)}(w)$ at $u = 2\lambda - 2e^\lambda \in (-\infty,-2)$ which by Lemma 2.5 is $2(e^\lambda - \lambda - 1)$. Furthermore, the lemma implies that there is $\eta = \eta(\lambda) > 0$ such that $\varphi_{sign(\lambda)}(w)$ is analytic on the slit disk $\{w : |w-u| < 2(e^\lambda-\lambda-1)(1+\eta)\}\setminus[-2,\infty)$. Since after appropriate translation and re-scaling this slit disk is larger than the indented disk $\Delta(\pi/4,\eta)$ introduced in [25, (2.5)], and the second part of Lemma 2.5 gives $\varphi_\pm(w) \approx -1/(w+2)$ as $w \to -2$, we can apply [25, Corollary 2] to get (1.11).

Finally, as $m_n(0) = 1$ the convergence at $\lambda = 0$ holds trivially. □

3. Proof of Theorem 1.3

We follow the method and notation of Dupuis-Ellis [20]. Although some arguments are analogous to those found in [20, Chapter 6] which considers random walk models with time-homogeneous continuous statistics, and [44] where a different model with a time singularity at $t = 0$ is examined, for completeness we give all relevant details as several differ, especially in the lower bound proof.

Let $X^n_j = Z_{j-k_0+1}/n$ for $k_0 \le j \le n$, and set $X^n_j = j/n$ for $0 \le j \le k_0$. Then, noting (1.1), given X^n_j, we have $X^n_{j+1} = X^n_j + v^n_j/n$ where v^n_j has Bernoulli distribution $\rho_{\sigma_n(j/n),X^n_j}$. Here,

$$\begin{aligned}\sigma_n(t) &= \begin{cases} s_{\lfloor nt-k_0+1\rfloor}/n & \text{for } t \ge \frac{k_0}{n}\\ 0 & \text{for } t < \frac{k_0}{n}\end{cases}\\ \rho_{\sigma,x}(A) &= \frac{x}{\sigma}\delta_0(A) + \Big(1-\frac{x}{\sigma}\Big)\delta_1(A) \qquad \text{for } A\subset\mathbb{R},\ 0\le x\le\sigma,\end{aligned}$$

and $\rho_{0,0} := \delta_1$.

Define $X^n_\cdot$ as the polygonal interpolated path connecting points $(j/n, X^n_j)$ for $0 \le j \le n$. Also, for probability measures $\mu \ll \nu$ such that $d\mu = f d\nu$, denote $R(\mu\|\nu) = \int f \log f d\nu$, the relative entropy; set $R(\mu\|\nu) = \infty$ when μ is not absolutely continuous with respect to ν.

Let $h : C([0,1],\mathbb{R}) \to \mathbb{R}$ be a bounded continuous function. To prove Theorem 1.3, we need only establish Laplace principle upper and lower bounds [20, page 74]. The upper bounds are to show

$$\limsup_{n\to\infty}\frac{1}{n}\log E\big\{\exp[-nh(X^n_\cdot)]\big\} \ \le\ -\inf_{\varphi\in\mathcal{C}_1}\big\{I(\varphi)+h(\varphi)\big\}$$

for a rate function I and a closed subset of continuous functions $\mathcal{C}_1$. The lower bounds are to prove the reverse inequality

$$\liminf_{n\to\infty}\frac{1}{n}\log E\big\{\exp[-nh(X^n_\cdot)]\big\} \ \ge\ -\inf_{\varphi\in\mathcal{C}_1}\big\{I(\varphi)+h(\varphi)\big\}.$$

Define, for $k_0+1 \le j \le n$, noting $X^n_j = j/n$ for $j \le k_0$ is deterministic, that

$$W^n(j,x_j,\dots,x_{k_0+1}) \ = \ -\frac{1}{n}\log E\Big[e^{-nh(X^n_\cdot)}\Big|X^n_j = x_j,\dots,X^n_{k_0+1} = x_{k_0+1}\Big],$$

and

$$W^n \ := \ W^n(k_0,\emptyset) \ = \ -\frac{1}{n}\log E\Big[e^{-nh(X^n_\cdot)}\Big].$$

Then, by the Markov property, for $k_0+1 \le j \le n-1$,

$$\begin{aligned}e^{-nW^n(j,x_j,\dots,x_{k_0+1})} &= E\Big[e^{-nW^n(j+1,X^n_{j+1},x_j,\dots,x_{k_0+1})}\Big|X^n_j = x_j,\dots,X^n_{k_0+1} = x_{k_0+1}\Big]\\ &= \int e^{-nW^n(j+1,x_j+v/n,x_j,\dots,x_{k_0+1})}d\rho_{\sigma_n(j/n),x_j}(dv).\end{aligned}$$

By a property of relative entropy [20, Prop. 1.4.2 (a)], for $k_0+1 \le j \le n-1$,

$$\begin{aligned}&W^n(j,x_j,\dots,x_{k_0+1})\\ &= -\frac{1}{n}\log\int e^{-nW^n(j+1,x_j+y/n,x_j,\dots,x_{k_0+1})}\rho_{\sigma_n(j/n),x_j}(dv)\\ &= \inf_\gamma\Big\{\frac{1}{n}R(\gamma\|\rho_{\sigma_n(j/n),x_j}) + \int W^n(j+1,x_j+y/n,x_j,\dots,x_1)\gamma(dy)\Big\}.\end{aligned}$$

Also, the boundary condition $W^n(n, x_n, \ldots, x_{k_0+1}) = h(x_\cdot)$ holds with respect to the linearly interpolated path $x_\cdot = x^n_\cdot$ connecting $\{(l/n, l/n)\}_{0\le l\le k_0}$,

The basic observation in the Dupuis-Ellis method is that $W^n(j, x_j, \ldots, x_{k_0+1})$ satisfies a control problem ([20, section 3.2]) whose solution for $k_0 \le j \le n-1$ is

$$V^n(j, x_j, \ldots, x_{k_0+1}) \;=\; \inf_{\{v_i^n\}} \bar{E}_{j,x_j,\ldots,x_{k_0+1}}\Big\{\frac{1}{n}\sum_{i=j}^{n-1} R(v_i^n(\cdot)\|\rho_{\sigma_n(i/n),\bar{X}_i^n}) + h(\bar{X}^n_\cdot)\Big\}.$$

Here, $v_i^n(dy) = v_i^n(dy; x_{k_0}, \ldots, x_i)$ is a Bernoulli distribution given $x_{k_0}, \ldots, x_i$ for $k_0 \le i \le n-1$ and in the display $\nu_i^n(\cdot) = \nu_i^n(dy|\bar{X}^n_{k_0}, \ldots, \bar{X}^n_i)$; $\{\bar{X}^n_i; 0 \le i \le n\}$ is the adapted path satisfying $\bar{X}^n_l = l/n$ for $0 \le l \le k_0$, and $\bar{X}^n_{l+1} = \bar{X}^n_l + \bar{Y}^n_l/n$ for $k_0 \le l \le n-1$ where $\bar{Y}^n_l$, conditional on $(\bar{X}^n_l, \ldots \bar{X}^n_{k_0})$ has distribution $v^n_l(\cdot)$; $\bar{X}^n_\cdot$ is the interpolated path with respect to $\{\bar{X}^n_l\}$; and $\bar{E}_{j,x_j,\ldots,x_{k_0+1}}$ denotes conditional expectation with respect to the $\bar{X}^n_\cdot$ process given the values $\{\bar{X}^n_l = x_l : k_0+1 \le l \le j\}$ at step $k_0+1 \le j \le n$. The boundary conditions are $V^n(n, x_n, \ldots, x_{k_0+1}) = h(x_\cdot)$ and

$$V^n(k_0, \emptyset) = V^n = \inf_{\{v_j^n\}} \bar{E}\Big\{\frac{1}{n}\sum_{j=k_0}^{n-1} R(v_j^n(\cdot)\|\rho_{\sigma_n(j/n),\bar{X}_j^n}) + h(\bar{X}^n_\cdot)\Big\}. \tag{3.1}$$

In particular, by [20, Corollary 5.2.1],

$$W^n \;=\; -\frac{1}{n}\log E\Big[e^{-nh(X^n_\cdot)}\Big] \;=\; V^n. \tag{3.2}$$

The goal will be to take Laplace limits using this representation. To simply later expressions, we will take $v_j^n = \delta_1$ for $0 \le j \le k_0 - 1$ when $k_0 \ge 1$.

3.1. **Upper bound.** To establish the upper bound, we first put the controls $\{v_j^n\}$ into continuous time paths: Let $v^n(dy|t) = v_j^n(dy)$ for $t \in (j/n, (j+1)/n]$, $0 \le j \le n-1$, and $v^n(dy|0) = v_0^n$. Define

$$v^n(A \times B) \;=\; \int_B v^n(A|t)dt$$

for $A \subset \mathbb{R}$, $B \subset [0,1]$. Define also the piecewise constant path $\tilde{X}^n(t) = \bar{X}^n_j$ for $t \in (j/n, (j+1)/n]$, $0 \le j \le n-1$, and $\tilde{X}^n(0) = 0$. Then,

$$V^n \;=\; \inf_{\{v_j^n\}} \bar{E}\Big\{\int_0^1 R(v^n(\cdot|t)\|\rho_{\sigma_n(\lfloor nt\rfloor/n),\tilde{X}^n(t)})dt + h(\bar{X}^n_\cdot)\Big\}. \tag{3.3}$$

From this control representation, as $|V^n| = |W^n| \le \|h\|_\infty$ and $\rho_{\sigma,x}$ is supported on $K = \{0,1\}$, for each n, there is $\{v_j^n\}$ supported on K and corresponding $v^n(dy \times dt) = v^n(dy|t) \times dt$ such that

$$W^n + \epsilon \;=\; V^n + \epsilon \;\ge\; \bar{E}\Big[\int_0^1 R(v^n(\cdot|t)\|\rho_{\sigma_n(\lfloor nt\rfloor/n),\tilde{X}^n(t)})dt + h(\bar{X}^n_\cdot)\Big].$$

As the sets K and

$$\Gamma \;=\; \Big\{\varphi \in C([0,1];\mathbb{R}) : \ \varphi(0) = 0, \ \varphi\uparrow, \text{ Lipschitz, with bound } 1\Big\}$$

are compact on $\mathbb{R}$ and $C([0,1],\mathbb{R})$ respectively, and $\{v_j^n\}$ are probability measures on K and $\{\bar{X}^n_\cdot\} \subset \Gamma$, by Prokhorov's theorem, the distributions of $(v^n(dy \times dt), \bar{X}^n_\cdot)$ have a subsequence which converges weakly to a limit distribution governing $(v, \bar{X}_\cdot)$ where for each realization, v is a probability measure on $K \times [0,1]$

and $\bar{X}_\cdot \in \Gamma$. More precisely, let $(\bar{\Omega}, \bar{F}, \bar{P})$ be a probability space where $v : \bar{\Omega} \to$ probability measures on $K \times [0,1]$, and $\bar{X} : \bar{\Omega} \to \Gamma$. Then, [20, Lemma 3.3.1] gives that v is the subsequential weak limit of v^n, and $\bar{P}$-a.s. for $\omega \in \bar{\Omega}$,

$$v(A \times B|\omega) \ = \ \int_B v(A|t,\omega)dt$$

for some kernel $v(dy|t,\omega)$ on K given t and ω.

Now, by the same proof given for [20, Theorem 5.3.5] (only [20, equation (5.12)] in the theorem statement differs; in our context μ there is replaced by $\rho_{\sigma_n(j/n),\tilde{X}^n_j}$), as K is compact, we have $(v^n, \bar{X}^n, \tilde{X}^n)$ has a subsequential limit in distribution to $(v, \bar{X}, \bar{X})$ [the last coordinate with respect to $D([0,1],\mathbb{R})$]. Also, $\bar{P}$-a.s., for all $t \in [0,1]$,

$$\bar{X}(t) \ = \ \int_{\mathbb{R}\times[0,1]} yv(dy \times ds) \ = \ \int_0^t \int_K yv(dy|s)ds.$$

In particular, $\bar{P}$-a.s., $\dot{\bar{X}}(t) = \int_K yv(dy|t)$.

By Skorohod representation theorem, we can assume $(v^n, \bar{X}^n, \tilde{X}^n) \to (v, \bar{X}, \bar{X})$ converges a.s. In particular, $\bar{X}^n$ converges uniformly to $\bar{X}$ a.s., and it is clear that $\tilde{X}^n$ converges uniformly to $\bar{X}$ a.s. as $\bar{X}$ is continuous ([20, p. 154]).

Let $\mu_\delta(\cdot) = (1-\delta)^{-1}1_{[\delta,1]}(\cdot)$ and $\nu^n(dy \times \mu_\delta dt) = \nu^n(dy|t) \times \mu_\delta dt$. Then, by [20, Lemma 1.4.3 (f)] we have, for $0 < \delta < 1$, that

$$\begin{aligned}&\int_\delta^1 R\big(v^n(\cdot|t)\big\|\rho_{\sigma_n(\lfloor nt\rfloor/n),\tilde{X}^n(t)}\big)dt\\ &\qquad = \ (1-\delta)R\big(v^n(dy \times \mu_\delta dt)\Big\|\rho_{\sigma_n(\lfloor nt\rfloor/n),\tilde{X}^n(t)} \times \mu_\delta dt\big).\end{aligned}$$

Write,

$$\begin{aligned}\liminf_{n\to\infty} V^n \ &\geq \ \liminf_{n\to\infty} \bar{E}\Big[\int_\delta^1 R\big(v^n(\cdot|t)\big\|\rho_{\sigma_n(\lfloor nt\rfloor/n),\tilde{X}^n(t)}\big)dt + h(\bar{X}^n_\cdot)\Big]\\ &\geq \ \liminf_{n\to\infty} \bar{E}\Big[(1-\delta)R\Big(v^n(dy \times \mu_\delta dt)\Big\|\rho_{\sigma_n(\lfloor nt\rfloor/n),\tilde{X}^n(t)} \times \mu_\delta dt\Big) + h(\bar{X}_\cdot)\Big]\\ &\geq \ \bar{E}\Big[\int_\delta^1 R(v(\cdot|t)\|\rho_{\alpha t,\bar{X}(t)})dt + h(\bar{X}_\cdot)\Big]\end{aligned}$$

where in the last line we use Fatou's lemma, noting lower semi-continuity of R, $\nu^n(dy \times \mu_\delta dt) \to \nu(dy|t) \times \mu_\delta dt$ a.s., and $\rho_{\sigma_n(\lfloor nt\rfloor/n),\tilde{X}^n(t)}$ converges in distribution to $\rho_{\alpha t,\bar{X}(t)}$ for $t \in [\delta,1]$ a.s. since $\sigma_n(\lfloor nt\rfloor/n) \to \alpha t$, $\tilde{X}^n(t) \to \bar{X}(t)$ uniformly on $[0,1]$ a.s., and $\rho_{\sigma,x}$ is continuous on $\{(\sigma,x) : \delta \leq \sigma \leq 1, 0 \leq x \leq \sigma\}$ (cf. Section 6.2 [20]).

By [20, Lemma 3.3.3(b)],

$$R(v(\cdot|t)\|\rho_{\alpha t,\bar{X}(t)}) \ \geq \ L\left(\alpha t, \bar{X}(t), \int yv(dy|t)\right)$$

where

$$L(t,x,y) \quad = \quad \sup_\theta \Big\{\theta y - \log \int e^{\theta z}\rho_{t,x}(dz)\Big\}. \tag{3.4}$$

We note, for $t > 0$, $L(t, x, y)$ diverges when $x = 0$, $0 \le y < 1$, and $x = t$, $0 < y \le 1$ but is finite when $0 < x < t$ and $0 \le y \le 1$, and in this case evaluates to

$$L(t, x, y) \;=\; y \log\left(\frac{ty}{t-x}\right) + (1-y)\log\left(\frac{t(1-y)}{x}\right), \tag{3.5}$$

understood with the usual convention $0 \log(0) = 0$.

Since $\int y v(dy|t) = \dot{\bar{X}}(t)$, we have

$$\liminf_{n\to\infty} V^n \;\ge\; \bar{E}\Big[\int_\delta^1 L(\alpha t, \bar{X}(t), \dot{\bar{X}}(t))dt + h(\bar{X}.)\Big].$$

As $L \ge 0$, and $\bar{X}(\cdot) \in \Gamma$, letting $\delta \downarrow 0$, we obtain,

$$\liminf_{n\to\infty} V^n \;\ge\; \inf_{\varphi\in\Gamma} \int_0^1 L(\alpha t, \varphi(t), \dot{\varphi}(t))dt + h(\varphi).$$

Taking into account (3.2), the upper bound holds with $\mathcal{C}_1 = \Gamma$ and

$$I(\varphi) \;=\; \int_0^1 L(\alpha t, \varphi(t), \dot{\varphi}(t))\, dt.$$

3.2. **Lower bound.** In the following, for typographical convenience, we write $E(X; \mathbb{A})$ for $\int_{\mathbb{A}} X dP$. Let now $\psi^* \in \Gamma$ be such that $I(\psi^*) < \infty$, and fix a bounded, continuous (in the sup norm) function $h : C([0, 1]; \mathbb{R}) \to \mathbb{R}$. In view of (3.2), we need only show, for each $\varepsilon > 0$, that

$$\limsup_{n\to\infty} V^n \;\le\; I(\psi^*) + h(\psi^*) + 8\varepsilon. \tag{3.6}$$

Step 1. Our first goal is to replace ψ^* by its appropriate regularization. We use the trick of considering a convex combination of paths,

$$\psi_\theta(t) \;=\; (1-\theta)\psi^*(t) + \theta t$$

for $0 \le \theta \le 1$. Since $\|\psi_\theta - \psi^*\|_\infty \le 2\theta$, it is clear, for small enough $\theta > 0$, that $|h(\psi_\theta) - h(\psi^*)| < \varepsilon$. Further, since I is finite on the line $y = t$ with slope 1, by convexity arguments [20, Lemma 1.4.3 (b)], for small enough $\theta > 0$,

$$I(\psi_\theta) \;\le\; I(\psi^*) + \varepsilon.$$

We therefore fix $\theta > 0$ such that $I(\psi_\theta) < I(\psi^*) + \varepsilon$ and $h(\psi_\theta) < h(\psi^*) + \varepsilon$.

Next, following [20, p. 82], we write

$$\varphi_\kappa(t) \;=\; \int_0^t \gamma_\kappa(s)ds, \tag{3.7}$$

where

$$\gamma_\kappa(t) \;=\; \kappa \int_{i/\kappa}^{i+1/\kappa} \dot{\psi}_\theta(s)ds$$

for $t \in (i/\kappa, (i+1)/\kappa]$, $0 \le i \le \kappa - 1$, and $\gamma_\kappa(0) = \gamma_\kappa(1/\kappa)$. For large enough κ, we have

$$\frac{\varphi_\kappa(t)}{t} \;\ge\; \theta \;>\; 0 \tag{3.8}$$

$$h(\varphi_\kappa) \;\le\; h(\psi^*) + 2\varepsilon \tag{3.9}$$

$$I(\varphi_\kappa) \;\le\; I(\psi^*) + 2\varepsilon. \tag{3.10}$$

Inequality (3.8) is a property of ψ_θ, and is preserved by (3.7). Since

$$\lim_{\kappa\to\infty}\sup_{t\in[0,1]}|\psi_\theta(t)-\varphi_\kappa(t)| \ = \ 0,$$

inequality (3.9) follows from continuity of h by our choice of θ. Then, as

$$\dot{\varphi}_\kappa(t) \ = \ \kappa\int_{\lfloor t\kappa\rfloor/\kappa}^{\lceil t\kappa\rceil/\kappa}\dot{\psi}_\theta(s)ds \ \to \ \dot{\psi}_\theta(t)$$

as $\kappa\uparrow\infty$, and $(t,x,y)\mapsto L(\alpha t,x,y)$ (cf. (3.4),(3.5)) as $\alpha>1$ is bounded, uniformly continuous on the compact set

$$\Big\{(t,x,y):\ 0\le t\le 1,\ \theta t\le x\le t,\ \theta\le y\le 1\Big\},$$

by the dominated convergence theorem, $\lim_\kappa I(\varphi_\kappa) = I(\psi_\theta)$. Inequality (3.10) follows due to our choice of θ.

We now fix κ such that the above bounds hold.

Since (3.10) implies $I(\varphi_\kappa)<\infty$, we now choose a $0<\delta<1/3$ such that

$$\int_0^\delta L(\alpha t,\varphi_\kappa(t),\dot{\varphi}_\kappa(t))dt \ < \ \varepsilon. \tag{3.11}$$

We will also need an estimate on s_n. From assumptions (1.2), there is an $0<\eta<1/2$ and $k_1\ge k_0+1$ such that for $n\ge k_1$,

$$\eta \ \le \ \frac{n}{s_{n-k_0+1}} \ \le \ 1-\eta. \tag{3.12}$$

With respect to η and θ, we impose additionally that δ satisfies

$$-\,\delta\log(\delta\eta\theta) \ < \ \varepsilon/10. \tag{3.13}$$

Step 2. We now build a sequence of controls based on φ_κ. Recall that we already set $v_j^n(dy)=\delta_1$ for $0\le j\le k_0-1$ when $k_0\ge 1$. Define

$$v_j^n(dy;x_{k_0},\ldots,x_j) \ = \ \begin{cases}\rho_{\sigma_n(j/n),x_j} & \text{for } k_0\le j\le k_1\\ \rho_{1,1-\dot{\varphi}_\kappa(j/n)} & \text{for } j\ge k_1+1.\end{cases}$$

Note, for $j\ge k_1+1$, ν_j^n does not depend on auxiliary inputs $x_{k_0},\ldots,x_j$, and is in fact the Bernoulli distribution with success probability $\dot{\varphi}_\kappa(j/n)$.

Define also $\bar{X}_l^n=l/n$ for $0\le l\le k_0$, and $\bar{X}_{j+1}^n=\bar{X}_j^n+\bar{Y}_j^n/n$ for $j\ge k_0$ where

$$\bar{P}(\bar{Y}_j^n\in dy|\bar{X}_0^n,\ldots,\bar{X}_j^n) \ = \ v_j(dy;\bar{X}_0^n,\ldots,\bar{X}_j^n).$$

Thus, for $j\ge 1$, $\bar{X}_j^n=(1/n)\sum_{\ell=0}^{j-1}Y_\ell^n$ where $\{Y_j^n\}_{j\ge k_1+1}$ are independent Bernoulli random variables with corresponding means $\{\dot{\varphi}_\kappa(j/n)\}_{j\ge k_1+1}$.

Step 3. We now collect some useful estimates.

A. Since $\bar{Y}_0^n\equiv 1$ (when $k_0=0$, recall $\rho_{\sigma,0}=\delta_1$), and the increments are at most one, we have $1/n\le\bar{X}_j^n\le j/n$ for $1\le j\le n$.

B. We have

$$\lim_{n\uparrow\infty}\sup_{0\le j/n\le 1}\left|\bar{X}_j^n-\frac{1}{n}\sum_{l=0}^{j}\dot{\varphi}_\kappa(l/n)\right| \ = \ 0 \qquad \text{a.s.} \tag{3.14}$$

Indeed, for large enough n, as $0 \le \dot\varphi_\kappa \le 1$, $(1/n)\sum_{j=0}^{k_1} |\dot\varphi_\kappa(j/n)| + k_1/n < n^{-1/8}$. Then, by Doob's maximal inequality,

$$\begin{aligned}
&\bar P\Big[\sup_{0\le j/n\le 1}\Big|\bar X_j^n - \frac{1}{n}\sum_{l=0}^{j}\dot\varphi_\kappa(l/n)\Big| > \frac{\theta}{2n^{1/8}}\Big] \\
&\quad\le\; Cn^{1/2}\bar E\Big|\bar X_n^n - \bar E(X_n^n)\Big|^4 \\
&\quad\le\; \frac{1}{n^{7/2}}\Big[C\sum_{l=0}^{n}\dot\varphi_\kappa(l/n)(1-\dot\varphi_\kappa(l/n))(1-3\dot\varphi_\kappa(l/n)+3\dot\varphi_\kappa^2(l/n)) \\
&\qquad\qquad + C\left(\sum_{l=0}^{n}\dot\varphi_\kappa(l/n)(1-\dot\varphi_\kappa(l/n))\right)^2\Big] \;\le\; Cn^{-3/2}. \qquad (3.15)
\end{aligned}$$

where C is a constant changing line to line.

C. For $0 \le j \le n$, from (3.8), it follows that

$$\frac{1}{n}\sum_{l=0}^{j}\dot\varphi_\kappa(l/n) \;\ge\; \theta\frac{j}{n}. \qquad (3.16)$$

D. Let $j \ge k_1 + 1$. Noting $1/n \le \bar X_j^n \le j/n$ (cf. part A) and bounds (3.12), we have

$$\eta \le 1 - \frac{j}{s_{j-k_0+1}} \le 1 - \frac{\bar X_j^n}{\sigma_n(j/n)} \le 1, \quad \frac{1}{s_{j-k_0+1}} \le \frac{\bar X_j^n}{\sigma_n(j/n)} \le 1-\eta.$$

Hence, $L(\sigma_n(j/n), \bar X_j^n, \dot\varphi_\kappa(j/n))$ can be well evaluated (cf. (3.4), (3.5)), and we may rewrite the relative entropy as

$$\begin{aligned}
R(v_j^n\|\rho_{\sigma_n(j/n),\bar X_j^n}) &= \dot\varphi_\kappa(j/n)\log[\dot\varphi_\kappa(j/n)/(1-\bar X_j^n/\sigma_n(j/n))] \\
&\quad +(1-\dot\varphi_\kappa(j/n))\log\big((1-\dot\varphi_\kappa(j/n))/(\bar X_j^n/\sigma_n(j/n))\big) \\
&= L(\sigma_n(j/n), \bar X_j^n, \dot\varphi_\kappa(j/n)).
\end{aligned}$$

Further, as $0 \le \dot\varphi_\kappa \le 1$, we have

$$\begin{aligned}
&L(\sigma_n(j/n), \bar X_j^n, \dot\varphi_\kappa(j/n)) \\
&\quad = \big[\dot\varphi_\kappa(j/n)\log\dot\varphi_\kappa(j/n) + (1-\dot\varphi_\kappa(j/n))\log(1-\dot\varphi_\kappa(j/n))\big] \\
&\qquad -\dot\varphi_\kappa(j/n)\log\big(1-\bar X_j^n/\sigma_n(j/n)\big) - (1-\dot\varphi_\kappa(j/n))\log\big(\bar X_j^n/\sigma_n(j/n)\big) \\
&\quad \le\; 0 - \log\eta + \log(s_{j-k_0+1}) \;\le\; \log(j/\eta^2), \qquad (3.17)
\end{aligned}$$

the last inequality using (3.12) again.

Step 4. We now argue (3.6) via representation (3.1). Let

$$\mathbb{A} \;=\; \Big\{\sup_{0\le j\le n-1}|\bar X_j^n - \frac{1}{n}\sum_{l=0}^{j}\dot\varphi_\kappa(l/n)| > \frac{\theta}{2n^{1/8}}\Big\}.$$

Since $R(v_j^n\|\rho_{\sigma_n(j/n),\bar{X}_j^n})=0$ for $k_0\le j\le k_1$, the sum in (3.1) equals

$$\begin{aligned}
\bar{E}\Big[\frac{1}{n}\sum_{j=k_0}^{n-1}R(v_j^n\|\rho_{\sigma_n(j/n),\bar{X}_j^n})\Big] &= \bar{E}\Big[\frac{1}{n}\sum_{j=k_1+1}^{n-1}L(\sigma_n(j/n),\bar{X}_j^n,\dot{\varphi}_\kappa(j/n))\Big]\\
&= \bar{E}\Big[\frac{1}{n}\sum_{j=k_1+1}^{n-1}L(\sigma_n(j/n),\bar{X}_j^n,\dot{\varphi}_\kappa(j/n));\mathbb{A}\Big]\\
&\quad + \bar{E}\Big[\frac{1}{n}\sum_{j=k_1+1}^{n-1}L(\sigma_n(j/n),\bar{X}_j^n,\dot{\varphi}_\kappa(j/n));\mathbb{A}^c\Big]\\
&= A_1+A_2. \qquad (3.18)
\end{aligned}$$

Step 5. We now treat the first term A_1 in (3.18). Combining (3.17) with (3.15), we obtain, for large n,

$$\begin{aligned}
&\bar{E}\Big[\frac{1}{n}\sum_{j=k_1+1}^{n-1}L(\sigma_n(j/n),\bar{X}_j^n,\dot{\varphi}_\kappa(j/n));\ \sup_{0\le j\le n-1}|\bar{X}_j^n-\frac{1}{n}\sum_{l=0}^{j}\dot{\varphi}_\kappa(l/n)|>\frac{\theta}{2n^{1/8}}\Big]\\
&\le\ \log(n/\eta^2)\bar{P}\Big[\sup_{0\le j/n\le 1}\Big|\bar{X}_j^n-\frac{1}{n}\sum_{l=0}^{j}\dot{\varphi}_\kappa(l/n)\Big|>\frac{\theta}{2n^{1/8}}\Big]\\
&\le\ C\log(n/\eta^2)n^{-3/2}\ <\ \varepsilon. \qquad (3.19)
\end{aligned}$$

Step 6. For the other term A_2 in (3.18), we split it into two sums depending on when index $j\le\delta n$ or $j\ge\delta n$ (recall δ from (3.11)):

$$\begin{aligned}
&\bar{E}\Big[\frac{1}{n}\sum_{j=k_1+1}^{\lfloor\delta n\rfloor}L(\sigma_n(j/n),\bar{X}_j^n,\dot{\varphi}_\kappa(j/n));\mathbb{A}^c\Big]\\
&\qquad + \bar{E}\Big[\frac{1}{n}\sum_{j=\lceil\delta n\rceil}^{n-1}L(\sigma_n(j/n),\bar{X}_j^n,\dot{\varphi}_\kappa(j/n));\mathbb{A}^c\Big]\\
&= B_1+B_2. \qquad (3.20)
\end{aligned}$$

Step 7. To estimate B_1, we divide it further into two terms corresponding to sums on indices $j\le n^{7/8}$ and $n^{7/8}\le j\le\delta n$:

$$\begin{aligned}
&\bar{E}\Big[\frac{1}{n}\sum_{j=k_1+1}^{\lfloor n^{7/8}\rfloor}L(\sigma_n(j/n),\bar{X}_j^n,\dot{\varphi}_\kappa(j/n));\mathbb{A}^c\Big]\\
&\qquad + \bar{E}\Big[\frac{1}{n}\sum_{j=\lceil n^{7/8}\rceil}^{\lfloor\delta n\rfloor}L(\sigma_n(j/n),\bar{X}_j^n,\dot{\varphi}_\kappa(j/n));\mathbb{A}^c\Big]\\
&= D_1+D_2.
\end{aligned}$$

The first term D_1, using (3.17), is bounded for all large n by

$$n^{-1/8}\log(n^{7/8}/\eta^2)\ <\ \varepsilon. \qquad (3.21)$$

The second term D_2, using again (3.17), is bounded in absolute value by

$$-\delta\log\eta + \frac{1}{n}\sum_{j=\lceil n^{7/8}\rceil}^{\lfloor\delta n\rfloor} |\log(s_{j-k_0+1}/n)|$$
$$-\bar{E}\Big[\frac{1}{n}\sum_{j=\lceil n^{7/8}\rceil}^{\lfloor\delta n\rfloor}\log\bar{X}^n_j;\ \sup_{0\le j\le n-1}|\bar{X}^n_j - \frac{1}{n}\sum_{l=0}^{j}\dot{\varphi}_\kappa(l/n)| \le \frac{\theta}{2n^{1/8}}\Big].$$

Now, note that (3.12) implies for $j \ge k_1+1$ that

$$\frac{j}{(1-\eta)n} \ \le\ \frac{s_{j-k_0+1}}{n} \ \le\ \frac{j}{\eta n}.$$

Then, as $0<\delta<1/3$ and $0<\eta<1/2$, we have for large n that

$$\begin{aligned}\frac{1}{n}\sum_{j=\lceil n^{7/8}\rceil}^{\lfloor\delta n\rfloor}|\log(s_{j-k_0+1}/n)| \ &\le\ 2\int_0^\delta \max\Big\{\Big|\log\Big(\frac{x}{1-\eta}\Big)\Big|,\Big|\log\Big(\frac{x}{\eta}\Big)\Big|\Big\}\,dx\\ &\le\ -6\delta\log\delta - 2\delta\log\eta.\end{aligned}$$

Also, noting (3.16), we have for large n that

$$\begin{aligned}&-\bar{E}\Big[\frac{1}{n}\sum_{j=\lceil n^{7/8}\rceil}^{\lfloor\delta n\rfloor}\log\bar{X}^n_j;\ \sup_{0\le j\le n-1}|\bar{X}^n_j - \frac{1}{n}\sum_{l=0}^{j}\dot{\varphi}_\kappa(l/n)| \le \frac{\theta}{2n^{1/8}}\Big]\\ &\le\ -\frac{1}{n}\sum_{j=\lceil n^{7/8}\rceil}^{\lfloor\delta n\rfloor}\log\Big[\frac{\theta j}{n} - \frac{\theta}{2n^{1/8}}\Big]\\ &\le\ -\frac{1}{n}\sum_{j=\lceil n^{7/8}\rceil}^{\lfloor\delta n\rfloor}\log\Big[\frac{\theta j}{2n}\Big]\ \le\ -2\int_0^\delta \log\frac{\theta x}{2}\,dx\ \le\ -4\delta\log\delta - 2\delta\log\theta.\end{aligned}$$

By combining these estimates, we have D_2 is bounded by a function of δ,η,θ which, given (3.13), can be made small:

$$D_2 \ \le\ -10\delta\log\delta - 3\delta\log\eta - 2\delta\log\theta \ <\ \varepsilon. \tag{3.22}$$

Step 8. We now estimate the second term B_2 in (3.20). Note, for $n > \delta^{-8}$, by (3.16) the event

$$\Big\{\sup_{0\le j\le n-1}|\bar{X}^n_j - \frac{1}{n}\sum_{l=0}^{j}\dot{\varphi}_\kappa(l/n)| \le \frac{\theta}{2n^{1/8}}\Big\}\ \subset\ \Big\{\inf_{j\ge\delta n}\bar{X}^n_j \ge \frac{\delta\theta}{2}\Big\}.$$

Hence, for large n,

$$B_2 \ \le\ \bar{E}\Big[\frac{1}{n}\sum_{j=\lceil\delta n\rceil}^{n-1} L(\sigma_n(j/n),\bar{X}^n_j,\dot{\varphi}_\kappa(j/n));\mathbb{A}^c\cap\Big\{\inf_{j\ge\delta n}\bar{X}^n_j \ge \frac{\delta\theta}{2}\Big\}\Big].$$

Note, from assumption (1.2) and $\bar{X}^n_j \le j/n$, that $\delta/(1-\eta)\le\sigma_n(j/n)\le 1/\eta$ and $\bar{X}^n_j \le \sigma_n(j/n)(1-\eta)$ when $\delta\le j/n\le 1$ for all large n. Also, $L(t,x,y)$ is continuous, and therefore also bounded and uniformly continuous on the compact set (cf. definition of L (3.4), (3.5)),

$$\Big\{(t,x,y):\ \frac{\delta}{1-\eta}\le t\le 1/\eta,\ \frac{\delta\theta}{4}\le x\le(1-\eta/2)t,\ 0\le y\le 1\Big\}. \tag{3.23}$$

Then,

$$\begin{aligned}
&\limsup_{n\to\infty} \bar{E}\Big[\frac{1}{n}\sum_{j=\lceil \delta n\rceil}^{n-1} L(\sigma_n(j/n), \bar{X}^n_j, \dot{\varphi}_\kappa(j/n)); \mathbb{A}^c \cap \Big\{\inf_{j\geq \delta n} \bar{X}^n_j \geq \frac{\delta\theta}{2}\Big\}\Big] \\
&\leq \limsup_{n\to\infty} \frac{1}{n}\sum_{j=\lceil \delta n\rceil}^{n-1} L\Big(\sigma_n(j/n), \frac{1}{n}\sum_{l=0}^{j}\dot{\varphi}_\kappa(l/n), \dot{\varphi}_\kappa(j/n)\Big). \qquad (3.24)
\end{aligned}$$

Further,

$$\lim_{n\to\infty}\sup_{1\leq j\leq n}\Big|\frac{1}{n}\sum_{l=0}^{j}\dot{\varphi}_\kappa(l/n) - \int_0^{j/n}\dot{\varphi}_\kappa(s)ds\Big| \;=\; 0, \tag{3.25}$$

as $\dot{\varphi}_k$ is piecewise constant and bounded. Then, given the bounds on $\sigma_n(j/n)$ above, $\theta j/n \leq (1/n)\sum_{l=0}^{j}\dot{\varphi}_k(l/n) \leq j/n$ (cf. (3.16)), and uniform continuity of L on the compact set (3.23), we may analogously bound (3.24) by

$$\lim_{n\to\infty}\frac{1}{n}\sum_{j=\lceil \delta n\rceil}^{n-1} L(\sigma_n(j/n), \varphi_\kappa(j/n), \dot{\varphi}_\kappa(j/n)) \;=\; \int_\delta^1 L(\alpha t, \varphi_\kappa(t), \dot{\varphi}_\kappa(t))dt. \tag{3.26}$$

Step 9. Finally, with respect to the second term in (3.1), by (3.14) and (3.25), in the sup topology, $\lim_{n\to\infty} h(\bar{X}^n_\cdot) = h(\varphi_\kappa(\cdot))$.

We now combine all bounds to conclude the proof of (3.6). By (3.1), and bounds (3.19), (3.21), (3.22), (3.26), we have

$$\begin{aligned}
\limsup_{n\to\infty} V^n \;&\leq\; \limsup_{n\to\infty}\bar{E}\Big[\frac{1}{n}\sum_{j=k_1+1}^{n-1} L(\sigma_n(j/n), \bar{X}^n_j, \dot{\varphi}_\kappa(j/n)) + h(\bar{X}^n_\cdot)\Big] \\
&\leq\; 3\varepsilon + \int_\delta^1 L(\alpha t, \varphi_\kappa(t), \dot{\varphi}_\kappa(t))\,dt + h(\varphi_k).
\end{aligned}$$

Then, by (3.11), (3.9), and (3.10), we obtain (3.6).

4. Proofs of Theorems 1.1 and 1.4

We first address the proof of Theorem 1.1, and later the proof of Theorem 1.4 in Subsection 4.3. Since $Z_n/n = \frac{n+k_0-1}{n}X_{n+k_0-1}(1)$, by contraction principle Theorem 1.3 implies Theorem 1.1 with the rate function I given in (1.14). As $x \mapsto \lambda x$ is a bounded continuous function on $[0,1]$, by Varadhan's Integral Lemma (see [18, Theorem 4.3.1], or [20, Theorem 1.3.4]), this implies that the limit (1.11) exists, and equals

$$\begin{aligned}
\Lambda(\lambda) \;&=\; \sup_{\varphi}\{\lambda\varphi(1) - I(\varphi)\} \\
&=\; \sup_{\varphi:\varphi(0)=0}\Big\{\int_0^1 \big[\lambda\dot{\varphi}(t) - \dot{\varphi}(t)\log\frac{\alpha t\dot{\varphi}(t)}{\alpha t - \varphi(t)} \\
&\qquad\qquad -(1-\dot{\varphi}(t))\log\frac{\alpha t(1-\dot{\varphi}(t))}{\varphi(t)}\big]\,dt\Big\}. \qquad (4.1)
\end{aligned}$$

Direct derivation of formula (1.12) or even formula (2.1) for $\alpha = 2$ from (4.1) seems quite challenging (cf. [44]). The Euler equations are:

$$\begin{aligned}\frac{\ddot{\varphi}}{\dot{\varphi}(1-\dot{\varphi})} &= \frac{\alpha}{\alpha t - \varphi} - \frac{1}{\varphi}\\ \varphi(0) &= 0\\ \frac{\dot{\varphi}(1)}{1-\dot{\varphi}(1)} &= \frac{\alpha - \varphi(1)}{\varphi(1)} e^{\lambda}.\end{aligned}$$

Numerical evidence suggests that the solutions of the Euler equations indeed give the correct answer.

4.1. **Proof of Theorem 1.1.** In this section we show that one can use Theorem 1.3 to set up the differential equation (1.7) which implies formula (2.1). Recall notation for the moment generating function. As we already noted, Theorem 1.3 implies that

$$\frac{1}{n}\log m_n(\lambda) \ \to\ \Lambda(\lambda) \tag{4.2}$$

with $\Lambda(\lambda)$ given by (4.1).

Formula (2.1) follows from the following additional fact.

Proposition 4.1. $\Lambda(\lambda)$, *as defined by* (4.2), *is differentiable and satisfies equation* (1.7).

It is straightforward to verify that (2.1) satisfies (1.7). The following argument shows the uniqueness: Suppose Λ_1, Λ_2 are two solutions with initial condition $\Lambda(0) = 0$. If $\Lambda_1(t) = \Lambda_2(t)$ for some $t > 0$, then they coincide for all $t > 0$. Therefore, we must have $\Lambda_1(t) > \Lambda_2(t)$ for all $t > 0$. By the mean value theorem, there is $t_0 > 0$ such that $\Lambda_1'(t_0) - \Lambda_2'(t_0) > 0$. But the equation gives $\Lambda_1'(t) - \Lambda_2'(t) = \alpha\frac{e^{\Lambda_1(t)} - e^{\Lambda_2(t)}}{1-e^t} < 0$ for all $t > 0$. Similarly, if $\Lambda_1(t) = \Lambda_2(t)$ for some $t < 0$, then they coincide for all $t < 0$. Therefore, we must have $\Lambda_1(t) > \Lambda_2(t)$ for all $t < 0$. By the mean value theorem, there is $t_0 < 0$ such that $\Lambda_1'(t_0) - \Lambda_2'(t_0) = \frac{\Lambda_1(t) - \Lambda_2(t)}{t} < 0$. But the equation gives $\Lambda_1'(t) - \Lambda_2'(t) = \alpha\frac{e^{\Lambda_1(t)} - e^{\Lambda_2(t)}}{1-e^t} > 0$ for all $t < 0$.

4.2. **Proof of Proposition 4.1.** As mentioned in the introduction, the differential equation is easy to derive heuristically from (1.4); the main technical difficulty is in justifying the convergence of various expressions.

To this end, the key ingredient is the control of the complex zeroes of $m_n(z)$, based on (the proof of) [10, Theorem 1]. Note that the assumptions in the following Proposition 4.2 include s_n satisfying (1.2), and $s_n = \alpha n$ for $\alpha = 1/2, 1$ with $k_0 \le s_1$.

Proposition 4.2. *Suppose* $m_n(z)$ *is defined by* (1.3) *with initial condition* $m_1(\lambda) = e^{k_0\lambda}$ *such that* $\{s_n\}$ *satisfies* $s_1 \ge k_0$, $s_n \ge \max\{k_0, 1\}$ *for* $n \ge 2$. *We also assume that there is* $n_0 \ge 1$ *such that* $s_{n_0} > k_0$ *when* $k_0 = 1, 2, \dots$ *and that there is* $n_1 \ge 2$ *such that* $s_{n_1} > 1$ *when* $k_0 = 0$. *Then* $m_n(z) \ne 0$ *in the strip* $|\Im(z)| < \pi$.

Proof. We note that $m_n(\lambda) = p_n(e^{\lambda})$ for a polynomial $p_n(u)$ with non-negative coefficients. Then, to prove the result, we repeat the proof of [10, Theorem 1] to deduce that all complex zeros of $p_n(u)$ are real, and non-negative for $n \ge 1$. [Some minor details differ from [10].]

Since $m_n'(\lambda) = up_n'(u)$, (1.3) gives

$$\begin{aligned} p_{n+1}(u) &= \frac{u(1-u)}{s_n} p_n'(u) + up_n(u) \\ &= \frac{u}{s_n}(1-u)^{s_n+1} \frac{d}{du}\left[(1-u)^{-s_n} p_n(u)\right]. \end{aligned} \tag{4.3}$$

Note that $p_1(u) = u^{k_0}$ and that $s_1 \geq k_0$. We first give the proof for the case when $s_1 > k_0 \geq 1$. To reach our conclusion we prove by induction the following statement.

> For $n \geq 2$, $p_n(u)$ is a polynomial of degree $n - 1 + k_0$ with a root of multiplicity k_0 at $u = 0$ and $n - 1 \geq 1$ strictly negative simple roots.

As $s_1 > k_0$, polynomial $p_2(u) = \frac{1}{s_1} u^{k_0}((s_1 - k_0)u + k_0)$ satisfies the inductive statement. Suppose for some $n \geq 2$, polynomial $p_n(u)$ is of degree $n - 1 + k_0$, has $n - 1$ simple negative roots $u_1 < u_2 < \cdots < u_{n-1}$ and a root of multiplicity k_0 at $u = 0$. Then, from the first equality in (4.3), $p_{n+1}(u)$ has also a root of multiplicity k_0 at $u = 0$. Clearly, $\{u_1, u_2, \ldots, u_{n-1}, 0\}$ are n distinct roots of the expression under the derivative on the right hand side of (4.3). Since $\{u_j\}$ are simple roots, the function must cross the real line, so by Rolle's theorem, $p_{n+1}(u)$ has $n - 1$ distinct roots interlaced between the roots of $p_n(u)$. This shows that $p_{n+1}(u)$ has $n + k_0 - 1$ real roots in the interval $(u_1, 0]$.

To end the proof, we want to show that the last $(n + k_0)$th root of $p_{n+1}(u)$ is located to the left of u_1, so that all negative roots of $p_{n+1}(u)$ must be simple. To see this, we follow again [10]: The first equation in (4.3) shows that $p_{n+1}(u_1)$ and $p_n'(u_1)$ have opposite signs, and $p_n'(u_1) \neq 0$ as u_1 is a simple root. So $p_{n+1}(u_1) \neq 0$. Since polynomials $p_{n+1}(u)$ and $p_n(u)$ have positive leading coefficients and their degrees differ by 1, their signs are opposite as $u \to -\infty$. Since u_1 is the smallest root, $p_n(u)$ has constant sign for $u < u_1$ which matches the sign of $p_{n+1}(u_1)$. Thus $p_{n+1}(u)$ must eventually cross the real line to the left of u_1. This shows that $p_{n+1}(u)$ has n simple strictly negative roots, and a root of multiplicity k_0 at $u = 0$, ending the induction proof.

Next, suppose $s_1 = s_2 = s_{n_0-1} = k_0 \geq 1$, but $s_{n_0} > k_0$ for some $n_0 \geq 2$. Then $p_1(u) = p_2(u) = \cdots = p_{n_0}(u) = u^{k_0}$ and the inductive proof goes through with minor modifications, starting with $p_{n_0+1} = \frac{1}{s_{n_0}} u^{k_0}((s_{n_0} - k_0)u + k_0)$ that replaces $p_2(u)$ in the previous argument.

Finally, if $k_0 = 0$, then $p_1(u) = 1$, $p_2(u) = u$. Choose first $n_1 \geq 2$ such that $s_{n_1} > 1$ but $s_n = 1$ for $2 \leq n < n_1$. Since in this case the value of s_1 is irrelevant, we get $p_2(u) = \cdots = p_{n_1}(u) = u$. The induction proof proceeds with minor modifications, starting with $p_{n_1+1}(u) = \frac{1}{s_{n_1}} u((s_{n_1} - 1)u + 1)$. □

Lemma 4.3. *Λ is differentiable and* $\frac{m_n'(\lambda)}{nm_n(\lambda)} \to \Lambda'(\lambda)$.

Proof. Recall that if $f(\zeta)$ is holomorphic in $|\zeta| \leq R$, then for $0 < r < R$

$$\max_{|\zeta| \leq r} |f(\zeta)| \leq \frac{R+r}{R-r} |f(0)| + \frac{2r}{R-r} \max_{|\zeta| \leq R} \Re f(\zeta) \tag{4.4}$$

(cf. [14, Theorem 6.31(ii)]).

By Proposition 4.2, for fixed $n \geq 1$, the function $m_n(z)$ is holomorphic and nonzero in the strip $|\Im z| < \pi$. Since $m_n(z)$ is a polynomial in e^z with nonnegative coefficients, $m_n(t) > 0$ for all $t \in \mathbb{R}$, $m_n(t)$ is increasing on $\mathbb{R}$, and $|m_n(z)| \leq m_n(|z|)$.

The function $m_n(z)$ has a holomorphic logarithm $L_n(z)$ on the strip $|\Im z| < \pi$. Because $m_n(t) > 0$ for $t \in \mathbb{R}$, we may assume that $L_n(t) = \log(m_n(t))$ for all $t \in \mathbb{R}$. For each $t \in \mathbb{R}$, we can apply (4.4) to $f(\zeta) = n^{-1} L_n(t + \zeta)$, $|\zeta| < \pi$, with $R = 2\pi/3$ and $r = \pi/3$. This gives

$$\begin{aligned} \max_{|z-t| \leq \pi/3} n^{-1}|L_n(z)| &\leq 3|n^{-1}L_n(t)| + 2n^{-1} \max_{|z-t|<\pi/3} |\Re L_n(\zeta)| \\ &= 3|n^{-1}\log(m_n(t))| + 2n^{-1} \max_{|z-t|<\pi/3} \log|m_n(z)| \\ &\leq 3|n^{-1}\log(m_n(t))| + 2\log(m_n(|t| + 2\pi/3)). \end{aligned}$$

Since $e^{-n|t|} \leq m_n(t) \leq e^{n|t|}$, $\{n^{-1}L_n(z) : n \geq 1\}$ is a normal family (i.e., a uniformly bounded family of holomorphic functions) in the disk $|z - t| \leq \pi/3$.

We now note that $\{n^{-1}L_n(t)\}$ converges for all real t; this holds true by (4.2) when $\alpha > 1$, or by Proposition 2.1 when $s_n = n$ or $s_n = n/2$. A version of Vitalli's theorem, see [22, p. 9], implies that $n^{-1}L_n(z) \to \Lambda(z)$ in the strip $|\Im z| \leq \pi/3$, the convergence is uniform in each disc $|z - t| \leq \pi/3$, the limit $\Lambda(z)$ is an analytic function of the argument z in that strip, and all derivatives of $n^{-1}L_n$ converge to the corresponding derivatives of Λ. In particular, the sequence $n^{-1}m_n'(\lambda)/m_n(\lambda) = n^{-1}L_n'(\lambda)$ converges to $\Lambda'(\lambda)$ for all real λ. □

Conclusion of Proof of Proposition 4.1. By Lemma 4.3 the right hand side of (1.4) converges, so the left hand side must converge too: $\frac{m_{n+1}(\lambda)}{m_n(\lambda)} \to \exp L(\lambda)$ for some L uniformly in a neighborhood of λ. Since the limit of ratios implies the same limit for n-th roots, we get $\frac{1}{n}\log m_n(\lambda) \to L(\lambda)$, which identifies $L(\lambda) = \Lambda(\lambda)$ as the pressure. From Lemma 4.3, the derivative $m_n'(\lambda)/(nm_n(\lambda)) \to \Lambda'(\lambda)$, so passing to the limit in (1.4) one obtains the differential equation for the pressure. □

4.3. **Proof of Theorem 1.4.** In the following, s_n satisfies assumptions (1.2), or $s_n = n$ or $s_n = n/2$. The LLN follows from the strict convexity of I in Theorem 1.1, and $\Lambda'(0) = \alpha/(1 + \alpha)$.

For the CLT, we recall [13, Proposition 2] in our context: When $\sup_n m_n(\epsilon)^{1/n} < \infty$ for some $\epsilon > 0$, $0 \notin \mathrm{closure}(\cup_{n\geq 1} Ze_n)$ where Ze_n is the zero set of $m_n(z) = E[\exp\{zZ_n\}]$, and Z_n/n satisfies an LDP, then $(Z_n - E[Z_n])/\sqrt{n}$ converges in distribution to $N(0, \sigma^2)$ where $\sigma^2 = \Lambda''(0)$.

To verify assumptions, note by Theorem 1.1 that the LDP for Z_n/n holds and $\lim_{n\to\infty} n^{-1}\log m_n(\lambda) = \Lambda(\lambda)$ for all $\lambda \in \mathbb{R}$, and by Proposition 4.2 that $m_n(z)$ has no zeroes in the strip $|\Im(z)| < \pi$ for $n \geq 1$. Finally, $\Lambda''(0) = \alpha^2/[(1 + \alpha)^2(2 + \alpha)]$ to finish the proof. □

5. Concluding remarks

We now comment on some possible extensions.

1. *Cases* $\alpha = 1/2, 1$. Although we prove a LDP for Z_n/n when s_n is linear with slopes $\alpha = 1/2, 1$ (Remark 1.2, Proposition 2.1), the proof of Theorem 1.1 for a path LDP with respect to $Z_{\lfloor nt \rfloor}/n$, especially the lower bound argument, does not cover these cases. The difficulty is in controlling boundary behavior as estimate (3.12) is not available. It would be interesting to look further into these issues.

2. *Higher order statistics.* With respect to random graph models, one might ask about LDP's for the vector $\mathbf{Z}_n^k/n = \langle Z_1(n), \ldots, Z_k(n)\rangle/n$ where the jth component $Z_j(n)$ counts the number of vertices with degree $j \leq k$ for $k \geq 2$. In principle, our method to analyze the leaves can be used to study $\mathbf{Z}_n^k/n$. Indeed, the Dupuis-Ellis type arguments given here for a path LDP for the leaves $Z_1(\lfloor nt\rfloor)/n$ (Theorem 1.1) would seem to extend to the vector-valued paths $\mathbf{Z}_{\lfloor nt\rfloor}^k/n$.

However, to calculate the pressure

$$\mathbf{\Lambda}_k(\lambda_1, \ldots, \lambda_k) \ = \ \lim_{n\to\infty} \frac{1}{n} \log E\Big[\exp\Big\{\sum_{i=1}^k \lambda_i Z_i(n)\Big\}\Big],$$

as in Theorem 1.3 for the leaves, the differential equation which now arises for $\mathbf{\Lambda}_k$, in place of the ODE for Λ (1.7), is a quasilinear PDE with $k \geq 2$ independent variables. These PDE's, although in principle implicitly solved by the method of characteristics, unfortunately do not seem to admit explicit solutions, at least to the extent found here with respect $Z_1(n)/n$, a reason why we have focused on detailed investigations on the leaves. It would be of interest to study better these higher order questions.

Acknowledgements. We thank Professor S.R.S. Varadhan for illuminating discussions.

References

[1] Albert, R. and Barabási, A.-L. (2002). Statistical mechanics of complex networks. *Rev. Modern Phys.* **74,** 47–97.

[2] Aldous, D. (2001). Stochastic models and descriptive statistics for phylogenetic trees, from Yule to today. *Statistical Science.* **16**, 23-34.

[3] Athreya, K. B., Ghosh, A. P. and Sethuraman, S. (2008). Growth of preferential attachment random graphs via continuous-time branching processes. *Proc. Indian Acad. Sci. (Math. Sci.)* **118,** 473–494.

[4] Bakhtin, Y. and Heitsch, C. (2008). Large deviations for random trees. *J. Stat. Phys.* **132,** 551–560.

[5] Barabási, A.-L. and Albert, R. (1999). Emergence of scaling in random networks. *Science* **286,** 509–512.

[6] Benassi, A. (1996). Arbres et grandes déviations. In *Trees (Versailles, 1995).* vol. 40 of *Progr. Probab.* Birkhäuser, Basel pp. 135–140.

[7] Blum, M.G.B., Francois, O. (2005). On statistical tests of phylogenetic tree imbalance: The Sackin and other indices revisited. *Math. Biosciences* **195**, 141-153.

[8] Blum, M.G.B., Francois, O., Janson, S. (2006). The mean, variance and limiting distribution of two statistics sensitive to phylogenetic tree balance. *Ann. Appl. Probab., 4* **16**, 2195-2214.

[9] Bollobás, B., Riordan, O., Spencer, J. and Tusnády, G. (2001). The degree sequence of a scale-free random graph process. *Random Structures Algorithms* **18,** 279–290.

[10] Bona, M. Real zeros and normal distribution for statistics on Stirling permutations defined by Gessel and Stanley 2007.

[11] Bonato, A. (2008). *A course on the web graph* vol. 89 of *Graduate Studies in Mathematics.* American Mathematical Society, Providence, RI.

[12] Broutin, N. and Devroye, L. (2006). Large deviations for the weighted height of an extended class of trees. *Algorithmica* **46,** 271–297.

[13] Bryc, W. (1993). A remark on the connection between the large deviation principle and the central limit theorem. *Stat. Probab. Lett.* **18,** 253–256.

[14] Burckel, R. B. (1979). *An introduction to classical complex analysis. Vol. 1* vol. 82 of *Pure and Applied Mathematics.* Academic Press Inc. [Harcourt Brace Jovanovich Publishers], New York.

[15] Chung, F. and Lu, L. (2006). *Complex graphs and networks* vol. 107 of *CBMS Regional Conference Series in Mathematics*. Published for the Conference Board of the Mathematical Sciences, Washington, DC.

[16] Cooper, C. and Frieze, A. (2003). A general model of web graphs. *Random Structures Algorithms* **22,** 311–335.

[17] Dembo, A., Mörters, P. and Sheffield, S. (2005). Large deviations of Markov chains indexed by random trees. *Ann. Inst. H. Poincaré Probab. Statist.* **41,** 971–996.

[18] Dembo, A. and Zeitouni, O. (1998). *Large deviations techniques and applications* second ed. vol. 38 of *Applications of Mathematics (New York)*. Springer-Verlag, New York.

[19] Dereich, S. and Morters, P. Random networks with sublinear preferential attachment: Degree evolutions 2008.

[20] Dupuis, P. and Ellis, R. S. (1997). *A weak convergence approach to the theory of large deviations*. Wiley Series in Probability and Statistics: Probability and Statistics. John Wiley & Sons Inc., New York. A Wiley-Interscience Publication.

[21] Dupuis, P., Nuzman, C. and Whiting, P. (2004). Large deviation asymptotics for occupancy problems. *Ann. Probab.* **32,** 2765–2818.

[22] Duren, P. L. (1983). *Univalent functions* vol. 259 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, New York.

[23] Durrett, R. (2006). *Random Graph Dynamics*. Cambridge U. Press.

[24] Flajolet, P., Gabarró, J. and Pekari, H. (2005). Analytic urns. *Ann. Probab.* **33,** 1200–1233.

[25] Flajolet, P. and Odlyzko, A. (1990). Singularity analysis of generating functions. *SIAM J. Discrete Math.* **3,** 216–240.

[26] Gessel, I. and Stanley, R. P. (1978). Stirling polynomials. *J. Combinatorial Theory Ser. A* **24,** 24–33.

[27] Jabbour-Hattab, J. (1999) Martingales et grandes deviations pour les arbres binaires de recherche. *C.R. Acad. Sci. Paris* **328**, Serie 1, 805-810.

[28] Janson, S. (2004). Functional limit theorems for multitype branching processes and generalized Pólya urns. *Stochastic Process. Appl.* **110,** 177–245.

[29] Janson, S. Plane recursive trees, Stirling permutations and an urn model 2008.

[30] Janson, S., Kuba, M. and Panholzer, A. Generalized Stirling permutations, families of increasing trees and urn models 2008.

[31] Katona, Z. and Móri, T. F. (2006). A new class of scale free random graphs. *Statist. Probab. Lett.* **76,** 1587–1593.

[32] Kober, H. (1952). *Dictionary of conformal representations*. Dover Publications Inc., New York, N. Y.

[33] Kotz, S. and Balakrishnan, N. (1997). Advances in urn models during the past two decades. In *Advances in combinatorial methods and applications to probability and statistics*. Stat. Ind. Technol. Birkhäuser Boston, Boston, MA pp. 203–257.

[34] McKenzie, A. and Steel, M. (2000). Distributions of cherries for two models of trees. *Math. Biosci.* **164,** 81–92.

[35] Mitzenmacher, M. (2006). A brief history of generative models for power law and lognormal distributions. *Internet Math.* **1** 226-251.

[36] Móri, T. F. (2002). On random trees. *Studia Sci. Math. Hungar.* **39,** 143–155.

[37] Najock, D. and Heyde, C. C. (1982). On the number of terminal vertices in certain random trees with an application to stemma construction in philology. *J. Appl. Probab.* **19,** 675–680.

[38] Olver, F. W. J. (1974). *Asymptotics and special functions*. Academic Press [A subsidiary of Harcourt Brace Jovanovich, Publishers], New York-London. Computer Science and Applied Mathematics.

[39] Rosenberg, N. (2006). The mean and variance of the numbers of r-pronged nodes and r-caterpillars in Yule-generated genealogical trees. *Ann. Combinatorics* **10**, 129-146.

[40] Rudas, A., Tóth, B. and Valkó, B. (2007). Random trees and general branching processes. *Random Struct. Algorithms* **31,** 186–202.

[41] Smythe, R. T. and Mahmoud, H. M. (1994). A survey of recursive trees. *Teor. Ĭmovīr. Mat. Stat.* 1–29.

[42] Szymański, J. (1987). On a nonuniform random recursive tree. In *Random graphs '85 (Poznań, 1985)*. vol. 144 of *North-Holland Math. Stud.* North-Holland, Amsterdam pp. 297–306.

[43] Yule, G.U. (1924). A mathematical theory of evolution, based on the conclusions of Dr. J.C. Willis. *Philos. Trans. Roy. Soc. London Ser. B.* **213**, 21-87.

[44] Zhang, J. X. and Dupuis, P. (2008). Large-deviation approximations for general occupancy models. *Combin. Probab. Comput.* **17,** 437–470.

Department of Mathematical Sciences, University of Cincinnati, 2855 Campus Way, PO Box 210025, Cincinnati, OH 45221-0025, USA
E-mail address: `Wlodzimierz.Bryc@UC.edu`

Department of Mathematical Sciences, University of Cincinnati, 2855 Campus Way, PO Box 210025, Cincinnati, OH 45221-0025, USA
E-mail address: `David.Minda@math.uc.edu`

Department of Mathematics, 396 Carver Hall, Iowa State University, Ames, IA 50011, USA
E-mail address: `sethuram@iastate.edu`